

DISSERTATIONES BIOLOGICAE UNIVERSITATIS TARTUENSIS

211

DISSERTATIONES BIOLOGICAE UNIVERSITATIS TARTUENSIS

211

PAULI SAAG

Natural variation
in plumage bacterial assemblages
in two wild breeding passerines



TARTU UNIVERSITY PRESS

Department of Zoology, Institute of Ecology and Earth Sciences,
Faculty of Science and Technology, University of Tartu, Estonia

Dissertation was accepted for the commencement of the degree of *Doctor philosophiae* in animal ecology at the University of Tartu on February 20, 2012 by the Scientific Council of the Institute of Ecology and Earth Sciences.

Supervisors: Prof. Raivo Mänd, University of Tartu, Estonia

Dr. Vallo Tilgar, University of Tartu, Estonia

Opponent: Prof. Edward H. Burt, Jr., Ohio Wesleyan University,
USA

Commencement: Room 301, 46 Vanemuise Street, Tartu, on May 4, 2012, at 10:15.

Publication of this thesis is granted by the Institute of Ecology and Earth Sciences, University of Tartu and by the Doctoral School of Earth Sciences and Ecology, created under the auspices of European Social Fund.



European Union
European Social Fund



Investing in your future

ISSN 1024–6479
ISBN 978–9949–19–963–1 (trükis)
ISBN 978–9949–19–964–8 (PDF)

Autoriõigus Pauli Saag, 2012

Tartu Ülikooli Kirjastus
www.tyk.ee
Tellimus nr 110

CONTENTS

LIST OF ORIGINAL PUBLICATIONS	6
1. INTRODUCTION.....	7
2. MATERIAL AND METHODS	12
2.1. Study system	12
2.2. Collecting field data and samples	13
2.3. Density measures of bacteria on feathers.....	14
2.4. Feather-degrading bacterial species richness	14
2.5. Feather color measures.....	15
2.6. Statistical analysis.....	15
3. RESULTS.....	16
3.1. Species differences.....	16
3.2. The effect of body topography.....	16
3.3. Habitat differences	16
3.4. Seasonal and annual changes	16
3.5. Sex differences.....	17
3.6. Bacteria, female condition and breeding output	17
3.7. Bacteria and plumage coloration.....	17
4. DISCUSSION	18
4.1. Species differences.....	18
4.2. The effect of body topography.....	19
4.3. Habitat differences	20
4.4. Seasonal changes.....	21
4.5. Annual variation.....	23
4.6. Sex differences.....	23
4.7. Bacteria, female condition and breeding parameters	24
4.8. Bacteria and plumage coloration.....	26
4.9. Conclusions.....	27
SUMMARY	28
SUMMARY IN ESTONIAN	30
REFERENCES.....	32
ACKNOWLEDGEMENTS	37
PUBLICATIONS.....	39

LIST OF ORIGINAL PUBLICATIONS

This thesis is a summary of the following papers, which are referred to in the text by the Roman numerals:

- I. Saag, P., Tilgar, V., Mänd, R., Kilgas, P., Mägi, M. 2011. Plumage bacterial assemblages in a breeding wild passerine: relationships with ecological factors and body condition. *Microbial Ecology*, 61: 740–749.
- II. Saag, P., Mänd, R., Tilgar, V., Kilgas, P., Mägi, M., Rasmann, E. 2011. Plumage bacterial load is related to species, sex, biometrics and fledging success in co-occurring cavity-breeding passerines. *Acta Ornithologica*, 46(2): 191–201.
- III. Saag, P., Kilgas, P., Mägi, M., Tilgar, V., Mänd, R. 2012. Inter-annual and body topographic consistency in the plumage bacterial load of Great tits. *Journal of Field Ornithology*, 83(1): 94–100.
- IV. Kilgas, P., Saag, P., Mägi, M., Tilgar, V., Mänd, R. Plumage bacterial load increases rapidly during nest-building in a passerine bird. *Journal of Ornithology*, in press, doi:10.1007/s10336–011–0801–3.
- V. Kilgas, P., Saag, P., Mägi, M., Edenberg, M., Tilgar, V., Mänd, R. Variation in assemblages of feather bacteria in relation to plumage coloration in female great tits (*Parus major*). *Condor*, accepted for publication.

Original publications are reproduced with the permission of the publishers.

I contributed to planning the work and collecting field data. I solely devised appropriate molecular and microbiological methods and performed all laboratory procedures and analyses, except plumage coloration analyses for paper V. I was primarily responsible for data analyses in papers I–III. I wrote the first drafts of papers I-III and contributed to the completion of all papers.

I. INTRODUCTION

Wild animals harbor a diverse community of bacteria and fungi (reviewed in Clayton 1999). Considering the important impact that symbiotic and parasitic microorganisms can have on their hosts (Hackstein & Van Alen 1996, Nuttall 1997), investigating microbe-host interactions may help to explain behavioral and reproductive variation within and between host populations. However, detailed research into the bacterial loads of wild animals and the influence of bacteria on hosts has been limited by the requirement for very complex methods. Recently, however, molecular and microbiological techniques have developed rapidly, and avian ecologists have adopted these methods with great enthusiasm (e.g. Burt & Ichida 1999, Lucas et al. 2005, Shawkey et al. 2005). The number of papers indexed in the Thomson Reuters Web of Science dealing with feather bacteria increased approximately seven-fold between the last decade of 20th century and the first decade of the new century, clearly illustrating the explosion of interest in the topic. The evidence generated by this recent work suggests that plumage bacteria play an important role in shaping the life histories of wild birds (Clayton & Moore 1997, Burt & Ichida 1999, Muza et al. 2000, Gunderson et al. 2009).

Birds make significant efforts to maintain plumage function and to control ectoparasite load. Preening and other forms of grooming are critical for limiting the abundance of feather lice and other arthropods (Hart 1997). Behaviors such as anting, dusting, sunning, and the inclusion of green vegetation in nesting material may also defend against ectoparasites and bacteria (Clayton 1999). Moreover, laboratory studies indicate that uropygial oil protects plumage – either chemically (Shawkey et al. 2003a, Ruiz-Rodriguez et al. 2009) or physically (Reneerkens et al. 2008) – against damage by feather-degrading bacilli (see also Møller et al. 2009).

Bird plumage can host various assemblages of bacteria and fungi, several of which are capable of degrading feather keratin (Sangali & Brandelli 2000, Lucas et al. 2003b, Riffel et al. 2003, Shawkey et al. 2003a). Bacterial damage to feathers may have important fitness consequences for wild birds. Plumage deterioration may result in decreased thermal insulation (Brush 1965) and aerodynamic efficiency (Swaddle et al. 1996). In the long-term, these negative effects might reduce parental survival and reproductive success; the latter via changes in parental condition (Burt & Ichida 1999, Muza et al. 2000) or indirectly through the trade-off between reproductive effort and self-preening behavior (Burt & Ichida 1999, Merilä & Hemborg 2000, Muza et al. 2000). Furthermore, feather-degrading bacteria could also affect plumage-based communication between birds by changing feather coloration. Given that plumage coloration is believed to reflect individual quality (Hamilton & Zuk 1982), bacterial-induced changes in this trait may reduce individual reproductive success by influencing social dominance and mate choice (Shawkey et al. 2007, Gunderson et al. 2009, Shawkey et al. 2009a, Ruiz.-de-Castaneda et al. 2012). At

the same time, many bacteria detected on feathers produce antimicrobial substances (Riley & Wertz 2002, Peralta-Sanchez et al. 2010) and may therefore play a role in protecting eggs from infections with pathogenic bacteria (Gunderson et al. 2008, Shawkey et al. 2009b, Peralta-Sanchez et al. 2010). However, certain feather-inhabiting bacteria may also act as pathogens (Bruce & Drysdale 1994). Little is known about interspecific antagonism and dominance within the bacterial communities inhabiting feathers, although such processes may play an important role in determining individual bird fitness. For example, when fast-growing generalist bacterial species (dominants) are present in a community, they might depress overall bacterial diversity (Martin-Platero et al. 2006, Soler et al. 2008, Peralta-Sanchez et al. 2010).

Feather-degrading bacteria probably associate with all species of wild birds (Burt 2009), and within-species many, if not all, individuals harbor such bacteria in their plumage (Gunderson et al. 2009, Peele et al. 2009). However, to date, relatively few studies have examined the factors that shape feather-degrading bacterial assemblages in avian plumage (see Muza et al. 2000, Shawkey et al. 2003a, Burt & Ichida 2004, Cristol et al. 2005, Lucas et al. 2005, Saranathan & Burt 2007). The variability of keratinolytic bacterial assemblages colonizing bird plumage is most likely related to avian feeding behavior (e.g., ground versus canopy birds – Burt & Ichida 1999) and soil characteristics (Lucas et al. 2003b). As bacterial communities differ between habitats depending on soil parameters, it has been suggested that the structure of bacterial assemblages in plumage is habitat dependent, and that this might result in differences even between individuals of the same bird species (Burt & Ichida 1999, 2004, Bisson et al. 2007). On the other hand, as characteristics of habitat might determine the exposure of birds to a particular set of bacteria, the evolution of habitat-specific defense mechanisms might be promoted (Ruiz-Rodriguez et al. 2009). Habitat differences and feeding behavior in combination with specific defense mechanisms against feather-degrading bacteria might also result in different bacterial loads on different body parts (Burt & Ichida 1999). However, there is shortage of published evidence concerning the habitat-dependent fitness consequences of feather degrading bacteria. Similarly, only a few studies have described seasonal changes in plumage bacteria assemblages (Burt & Ichida 1999), e.g. during the pre-breeding and breeding periods (Bisson et al. 2009).

To summarize the state of current knowledge, it appears fair to conclude that while many valuable studies have been conducted (Table 1), there remains a shortage of information about many aspects of feather bacteria ecology. The largest gaps in our knowledge are in understanding how plumage bacterial assemblages are related to a bird's sex, body condition and breeding effort, as well as to habitat and season. Such knowledge is crucial for planning experiments to determine and understand the causality of previously reported relationships: whether and how feather-bacteria influence a bird's body condition, breeding success and survival. In order to generalize current findings, more comparable data from different study systems and bird species are required.

Table 1. Previously described relationships between ecological and life-history traits and feather-degrading (FDB) and other plumage-inhabiting bacteria in birds.

Source of variation	Studied bacterial parameter	Effect direction	Reference
<i>Inter-individual differences</i>			
Sex	FDB abundance	various effects	Gunderson et al. <i>J. Avian Biol.</i> 2009
	FDB and/or total bacterial abundance	females > males	Lucas et al. <i>Mol. Ecol.</i> 2005; Møller et al. <i>Fun. Ecol.</i> 2009; Czirjak et al. <i>Microb. Ecol.</i> 2010
Brood size	Total bacterial abundance	positive correlation	Lucas et al. <i>Mol. Ecol.</i> 2006
	FDB diversity	positive correlation	Lucas et al. <i>Mol. Ecol.</i> 2005
	Total bacterial diversity	effect on FDB assemblage structure	Lucas et al. <i>Mol. Ecol.</i> 2005
Body mass	FDB abundance	various effects	Gunderson et al. <i>J. Avian Biol.</i> 2009
Preening gland size	FDB abundance	positive correlation	Møller et al. <i>Fun. Ecol.</i> 2009
<i>Intra-individual differences</i>			
Body parts	FDB abundance	venter > dorsum	Burt & Ichida <i>Auk</i> 1999
Feather regions	Total bacterial abundance	distal > proximal	Muza et al. <i>Wilson Bull.</i> 2000
Feather color	Feather resistance to FDB	no color effect	Mahler et al. <i>Ibis</i> 2010
	Feather resistance to FDB	black > white	Goldstein et al. <i>Auk</i> 2004; Gunderson et al. <i>J. Avian Biol.</i> 2008; Ruiz-de-Castaneda et al. <i>Biol. J. Linn. Soc.</i> 2012
	Feather resistance to FDB	light > dark	Grande et al. <i>Ardeola</i> 2004
	FDB abundance	color effect	Gunderson et al. <i>J. Avian Biol.</i> 2009; Burt et al. <i>Biol. Lett.</i> 2011
Feather hue	FDB abundance	negative correlation	Shawkey et al. <i>Naturwissenschaften</i> 2008
Feather chroma	FDB abundance	positive correlation	Shawkey et al. <i>Am. Nat.</i> 2007

Table 1 (Continued). Previously described relationships between ecological and life-history traits and feather-degrading (FDB) and other plumage-inhabiting bacteria in birds.

Source of variation	Studied bacterial parameter	Effect direction	Reference
<i>Behavioral activities</i>			
Anting	Total bacterial abundance	no effect	Revis & Waller <i>Auk</i> 2004
Molt	Total bacterial abundance	no effect	Giraudeau et al. <i>J. Avian Biol.</i> 2010
Preening	FDB abundance	negative effect on FDB growth	Shawkey et al. <i>J. Avian Biol.</i> 2003
Foraging habit	FDB abundance	ground feeding > canopy birds	Burt & Ichida <i>Auk</i> 1999
Migration	Total bacterial diversity	migratory > sedentary	Bisson et al. <i>Microb. Ecol.</i> 2009
<i>Environmental variables</i>			
Sunlight intensity	Total bacterial abundance	negative correlation	Saranathan & Burt <i>Wilson Journal of Ornithology</i> 2007
Habitat	FDB diversity	various effects	Bisson et al. <i>Microb. Ecol.</i> 2007
	FDB abundance	salt marsh > freshwater marsh	Peele et al. <i>Auk</i> 2009
	FDB activity	humid > arid	Burt & Ichida <i>Condor</i> 2004
Season	FDB abundance	winter > summer	Burt & Ichida <i>Auk</i> 1999
<i>Other Factors</i>			
Breeding pair	FDB and/or total bacterial abundance	intrapair correlation	Lucas et al. <i>Mol. Ecol.</i> 2005; Gunderson et al. <i>J. Avian Biol.</i> 2009
Population differences	FDB and total bacterial abundance	positive correlation with breeding colony size	Møller et al. <i>Fun. Ecol.</i> 2009; Czirjak et al. <i>Microb. Ecol.</i> 2010
Species differences	Total bacterial diversity	various effects	Shawkey et al. <i>Waterbirds</i> 2006
Antibacterial substances (AS) in preening oil	FDB abundance and keratinolysis	AS inhibits FDB and keratinolysis, produced by symbiotic bacteria	Martin-Platero et al. <i>Appl. Environ. Microbiol.</i> 2006; Martin-Vivaldi et al. <i>J. Avian Biol.</i> 2009; Ruiz-Rodriguez et al. <i>J. Exp. Biol.</i> 2009; Martin-Vivaldi et al. <i>Proc. Roy. Soc.</i> 2010
Captivity	FDB activity	no effect	Cristol et al. <i>Auk</i> 2005

The general aim of my thesis was to explore the extent, pattern and correlates of natural variation in numbers and species richness of bacterial assemblages inhabiting the plumage of two co-occurring cavity-breeding passerines, Great Tits (*Parus major*) and Pied Flycatchers (*Ficedula hypoleuca*), which share habitat and nesting site requirements, but differ markedly in certain other life-history traits. Specifically, I studied the following questions:

(i) Do plumage bacterial assemblages differ (both in terms of density and species richness) between these two bird species breeding in the same area and using the same habitats and nest sites? Similar bacterial patterns in comparable bird species would suggest that the factors shaping bacterial communities (e.g., sex, habitat, season, parental effort etc.) act in similar ways in different bird species. On the other hand, inter-specific differences would indicate that some bacterial characteristics are species-specific;

(ii) Do plumage bacterial assemblages differ between body regions in the studied bird species? I expected that more bacteria would be recorded on the ventral than the dorsal regions, because the ventral part usually comes into greater contact with potential contamination sources, including the ground (Burt & Ichida 1999). I also expected intra-individual differences in plumage bacterial load between different body parts to be consistent through time;

(iii) Do plumage bacterial assemblages differ between habitats? I expected that bacterial load would be higher in more heterogeneous (e.g., deciduous forest) than more homogeneous habitats (e.g., coniferous forest);

(iv) Do plumage bacterial assemblages vary through the breeding season? I expected bacterial load to increase during the breeding season due to the seasonal increase in air temperature (Burt & Ichida 1999, Peele et al. 2009), the increasing time of exposure to contamination sources, and potentially due to the cumulative effects of reproductive effort on individual condition (Lucas et al. 2005);

(v) Do plumage bacterial assemblages vary between years? I expected that different climatic conditions in different years would influence bacterial abundance to some extent. At the same time, if bacterial abundance reflects individual quality (e.g., resistance to bacterial infestation), consistent differences between individuals might be expected despite annual variability.

(vi) Do plumage bacterial assemblages differ between sexes? I expected females to harbor relatively higher loads of bacteria on their plumage than males, because of the more diverse reproductive activities undertaken by females (Lucas et al. 2005);

(vii) Are plumage bacterial assemblages related to parental body condition and reproductive output? I expected both relationships to be negative, due to possible negative influence that plumage bacteria may exert on host (see above);

(viii) Are bacterial assemblages related to plumage coloration? I expected high bacterial load to be associated with less bright/colorful plumage, because bacteria may damage feather structure (Shawkey et al. 2007, Gunderson et al. 2009, Shawkey et al. 2009a, Ruiz-de-Castaneda et al. 2012).

2. MATERIAL AND METHODS

2.1. Study system

The Great Tit (*Parus major*) is a small (18–19 g) insectivorous passerine, which is common throughout the palearctic region (Cramp & Perrins 1993). Great Tits mainly forage in the tree canopy during the breeding period, feeding mostly on Lepidoptera and sawfly larvae and spiders (Cramp & Perrins 1993, Gosler 1993). The species is only partly migratory in the study region (Vilbaste 1994), and if it migrates, then it does not usually travel very long distances. Many individuals spend the winter in the vicinity of their breeding area, usually close to human settlements (Perrins 1979, Gosler 1993, Vilbaste 1994). Great Tits inhabit various types of woodland, but prefer deciduous forests for breeding (Mänd et al. 2005). The species is a facultative double brooder, and in the study area 40–70% of females lay a second clutch during the breeding season (Mägi & Mänd 2004). In the study area, nest building starts at the end of April. The first breeding period lasts approximately from the end of April to the middle of June, while the second breeding period lasts from the end of June to the end of July. First clutches usually contain 9–12 eggs (Hörak et al. 1995), while second clutches are generally smaller (Mägi & Mänd 2004).

The Pied Flycatcher (*Ficedula hypoleuca*) is a small (12–13 g) insectivorous migratory passerine, that occurs throughout much of northern and eastern Europe (Lundberg & Alatalo 1992). Pied Flycatcher diet consists of various arthropods, and birds forage both in trees and on the ground (Lundberg & Alatalo 1992). Pied Flycatchers share the same habitats as Great Tits. Unlike Great Tits however, Pied Flycatchers do not lay two clutches. In the study area Flycatchers start to breed at least two weeks later than Great Tits, with breeding lasting from the end of May to the end of June. Pied Flycatcher clutches normally contain 6–7 eggs.

Besides sharing habitat preferences, both species breed in tree-holes and will also readily accept nest-boxes. Both species are also short-lived, with more than half of individuals breeding only once (Perrins 1979, Lundberg & Alatalo 1992). Hence, there are many similarities in the ecology of the studied species, but also several differences (see above).

All field studies were conducted in northern Europe, near Kilingi-Nõmme (58° 7'N, 25° 5'E), SW Estonia, in 2007 (only Great Tit) and 2008 (both species). The study area covers approximately 50 km² and contains a mosaic of two forest types – coniferous and deciduous (Figure 1).

Birds bred in wooden nest boxes with a cavity of 11 × 11 × 30 cm and an entrance diameter of 3.5 – 4.0 cm. Nest boxes were mounted on tree trunks at a height of 1.5 – 2.0 m and were distributed in both (coniferous and deciduous) habitats. The distance between neighboring nest boxes was 50–60 m. Nest boxes were cleaned to remove old nest material before the beginning of the breeding season.




Figure 1. Schematic map of the study area.

2.2. Collecting field data and samples

All nest boxes were checked to record the laying date of the first egg, the clutch size and the hatching dates of both the first and second broods (in Great Tits). These two brood categories were clearly distinguishable since there is no overlap between the laying dates of the first and second clutches. Second broods were also confirmed by ringing data, as at least one adult from each breeding pair had already been captured in a particular year (see below). The number of fledglings per nest was recorded. Adults were captured using automatic traps at their nests during the second half of the nestling period. In 2007 (but not in 2008), female Great Tits were also captured during the pre-laying (nest-building) stage (at night when roosting in nest boxes, see paper **IV** for details). Male Great Tits were more distrustful of the traps, compared with females; therefore, the male sample size was much smaller and unequally distributed between breeding attempts. In the case of Pied Flycatchers, there was no significant difference between sexes in terms of trapping efficiency. Trapped adults were individually marked with numbered metal rings. Birds were weighed with a Pesola spring balance to a precision of 0.1 g, and tarsus measurements taken to the nearest 0.1 mm and wing length to the nearest 1 mm using digital callipers.

A fresh pair of examination gloves was used each time a new bird was handled. Within 30 sec after capture, about 5 ventral feathers were plucked from the right side of each bird's chest, using forceps cleaned in 96% ethanol, and placed in dry clean disposable microtubes to assess the number of bacteria on feathers. In 2008, but not in 2007, the same number of dorsal feathers (from the center of the back) was also collected from Great Tits. Feathers from the different body regions were placed in separate tubes. In both years, another sample of five feathers (from the same body regions) was collected from each bird to determine the species richness of feather-degrading bacterial assemblages. See papers **I**, **III** and **IV** for details.

2.3. Density measures of bacteria on feathers

Two distinguishable ecological types of bacteria occur in bird plumages: free-living and attached bacteria. Studies of bacterial communities in soil, water and sediment have demonstrated that free-living bacteria are usually more labile, while attachment provides a more stable environment and protection against grazing, chemical antibiotics or physical stress (see Ozawa & Yamaguchi 1986, Lucas et al. 2003a, Selje & Simon 2003 for references).

Free-living bacteria were washed out from the feathers using PBS (phosphate buffered saline) solution. To remove attached bacteria, feathers were sonicated in detergent solution in order to break down biofilms (Lucas et al. 2003a). Free-living and attached bacteria samples were stored and analyzed separately. Direct counts were performed with a flow cytometry machine (BD LSR II) that was calibrated to detect only bacterium sized tagged particles. DNA-binding dye SYBR Green was used for tagging. The number of feathers in each sample was recorded in order to calculate bacterial density per feather (as Lucas et al. 2005). See paper **I** for details.

2.4. Feather-degrading bacterial species richness

At the laboratory, feather-degrading bacterial assemblages were enriched (following Lucas et al. 2005). The ribosomal intergenic spacer analysis (RISA) method was used to analyze the structure of feather-degrading assemblages obtained in the enrichment cultures. Each RISA band is assumed to correspond to one bacterial species and is referred to as a phylotype (following Muyzer et al. 1993, Stach et al. 2003, to point out that RISA bands are anonymous). Thus, the band profile reflects bacterial assemblage structure, while the number of bands corresponds to the bacterial assemblage richness (Ranjard et al. 2000). In this thesis, term phylotype is used for referring RISA results, otherwise term species is used.

This method is fairly coarse (and therefore easily applicable and cheap) and does not allow ecological characteristics (free-living or attached) to be assigned

to particular bacterial phylotypes. Also, due to the specific limitations of RISA analysis, a certain portion of true species richness might be overlooked. However, general estimates of diversity should still be reliable (Ranjard et al. 2000). See paper **I** for details.

2.5. Feather color measures

Three additional feathers were plucked from each Great Tit individual from a standard position in the yellow breast plumage. Breast color was characterized by measuring chroma values using a spectrophotometer (Ocean Optics USB2000 with Ocean Optics DH2000 lamp). Chroma corresponds to color purity on a scale from 0 to 100, with 100 representing pure color. Chroma was measured in the visible range of 400–700 nm (Senar et al. 2008; Broggi and Senar 2009). The three feathers were put on top of each other and three measurements were obtained from each individual. The mean was used in analyses. Within-individual repeatability of the three chroma measurements was high ($r = 0.86$; according to Lessells & Boag 1987). See paper **V** for details.

2.6. Statistical analysis

Statistical analysis was performed using program Statistica (Statsoft, Tulsa, Oklahoma). Bacterial density estimates were log-transformed prior to analysis to satisfy the assumption of normality. General linear models (GLMs) were used to model variation in bacterial density. Akaike information criterion (AIC) was used for model selection in study **II**, otherwise predictors were removed from the initial models using a backward stepwise procedure, when non-significant ($P > 0.05$). See original papers for more details.

3. RESULTS

3.1. Species differences

The density of attached bacteria on feathers was lower in Pied Flycatchers than Great Tits (**II**). Free-living bacterial density and the mean number of feather-degrading bacterial phylotypes per bird did not differ significantly between the two bird species. In total, 16 bacterial phylotypes were detected from 110 Great Tit individuals and 12 bacterial phylotypes were found from 42 Pied Flycatchers (**II**).

3.2. The effect of body topography

In Great Tits, the densities of free-living, but not attached, bacteria on dorsal and ventral feathers were positively correlated. Correlation between the richness of feather-degrading bacterial communities in these body areas was also positive, but marginally non-significant (**III**). At the same time, the densities of both attached and free-living bacteria were significantly higher on dorsal than on ventral feathers in Great Tits, while the richness of feather-degrading bacterial communities on dorsal and ventral feathers did not differ significantly (**III**).

3.3. Habitat differences

In Great Tits, the densities of both free-living and attached bacteria were significantly higher in individuals breeding in deciduous habitat compared with those breeding in coniferous habitat (**I, IV**). A habitat difference in the number of phylotypes was only apparent during the nest-building stage – individuals carried fewer phylotypes in deciduous than in coniferous habitat (**I**). By contrast, no significant effect of habitat type on the plumage bacterial community was detected in the Pied Flycatcher (**II**).

3.4. Seasonal and annual changes

The density of both free-living and attached bacteria on Great Tit feathers was higher during the nest-building period than during either the first-brood or second-brood stage (**I**). Moreover, the precise stage of nest-building also had a significant effect on the density of attached bacteria on Great Tit feathers, with the density tending to increase towards the final stage of nest-building (**IV**). The decline in bacterial density between the nest-building stage and the first brood was confirmed by repeated measures analysis of the same individuals in the case of attached but not free-living bacteria (**I**). Later in the season, the density of attached bacteria increased significantly from the first to second brood (**I**).

The density of attached bacteria on feathers was generally higher in 2008 than in 2007 (**I**, **III**). In six females, captured in the two consecutive years, there was a significant positive correlation between the densities of attached bacteria in the different years (**III**).

3.5. Sex differences

Males of both Great Tits and Pied Flycatchers combined had fewer attached bacteria on their plumage than females (**II**). The difference remained when data from Pied Flycatchers were analyzed separately, and males also had fewer bacterial phylotypes on their feathers than females on average (**II**). Male Great Tits, however, supported on average more bacterial phylotypes than females, and the difference was more pronounced during second broods than during first broods (**I**). No significant sex differences were found in the density of free-living bacteria. Both free-living and attached bacterial densities were correlated within breeding pair members.

3.6. Bacteria, female condition and breeding output

In Great Tits, a positive relationship between free-living bacterial density and female mass was found during both the first and second breeding periods (**I**), while the density of attached bacteria was negatively related to female mass only during the first breeding period (**I**). No significant effect of body mass was found on feather-degrading bacterial phylotypic richness. In Pied Flycatchers, there was a significant positive correlation between the density of attached bacteria and the tarsus length of adult birds, and a negative correlation between the number of bacterial phylotypes and body mass (**II**). No significant effect of body mass was found on free-living bacterial density.

There was a negative relationship between the density of attached bacteria and the number of fledglings in Great Tits but not in Pied Flycatchers (**II**). There was no such relationship with clutch size (**II**).

3.7. Bacteria and plumage coloration

In Great Tits, plumage chroma during the nestling-feeding period was negatively correlated with the phylotypic richness of feather-degrading bacteria, but not with densities of attached or free-living bacteria (**V**). The change in chroma between the pre-laying and chick-rearing periods was correlated with the change in attached bacterial density during same time period. This means that in individuals whose attached bacterial densities increased over time, chroma simultaneously decreased (**V**).

4. DISCUSSION

4.1. Species differences

The total number of bacterial phylotypes observed in Great Tits (18 phylotypes from 290 individuals, **I**) and Pied Flycatchers (12 phylotypes from 42 individuals, **II**) was of approximately the same magnitude as recorded in other studied wild bird species: in House Finches *Carpodacus mexicanus* (13 phylotypes from 29 individuals) and in Eastern Bluebirds *Sialia sialis* (15 phylotypes from 4 individuals) (Shawkey et al. 2003a, Shawkey et al. 2005). However, the mean richness of bacterial phylotypes per individual (the sample consisting of five ventral feathers) was relatively low both in Great Tits (2.3 ± 1.6 , **I**) and Pied Flycatchers (2.0 ± 1.0 , **II**), compared with that recorded from European Starlings *Sturnus vulgaris* (7 ± 3) using exactly the same method (Lucas et al. 2005). Given the relatively high abundance of feather-degrading bacteria in soil (Lucas et al. 2003b), differences in foraging behavior could be a possible explanation for such interspecific differences in bacterial diversity. Great Tits and Pied Flycatchers mainly forage in the canopy, while starlings are mainly ground-foragers. Similarly, other authors have found feather-degrading bacteria to be more abundant in the plumage of ground-feeding birds compared with canopy birds (Burt & Ichida 1999). However, one must consider that climatic and local soil or habitat parameters might also influence microbial communities, making it more difficult to compare different studies.

While the mean richness of bacterial phylotypes did not differ between the studied species, there was a significant difference in the density of attached bacteria: Pied Flycatchers harbored significantly fewer bacteria than Great Tits. However, on average Pied Flycatchers spend more time foraging on the ground than Great Tits (Haartman 1954, Alatalo & Alatalo 1979). Hence, the explanation of interspecific differences based on different foraging strategies does not seem to apply in a comparison of these two species. A possible alternative explanation is that Great Tit nests are much larger than those of Pied Flycatchers, contain more materials collected from the ground, and probably also harbor more feather-degrading bacteria. Also, it has previously been noted that migratory birds generally carry fewer bacteria on their feathers than sedentary birds (Bisson et al. 2009). The finding that the Pied Flycatcher, which is a long-distant migrant, has a lower average bacterial load than the mostly sedentary Great Tit, corresponds well with this general pattern. A third possible reason for the observed inter-specific differences in bacterial load is that the plumage coloration of these two species is very different, and feather color may influence the associated bacterial assemblage (Goldstein et al. 2004, Gunderson et al. 2008, Peralta-Sanchez et al. 2010, Peralta-Sanchez et al. 2011, see also Ruiz-de-Castaneda et al. 2012). However, the reasons underlying bacterial community differences seem to go beyond coloration, because Goodenough and Stallwood (2010) recently found that bacterial loads in nests differ significantly

even between bird species with as close biology and plumage coloration as Blue and Great Tits (*Cyanistes caeruleus et Parus major*), breeding in the same area.

Although it is impossible to draw firm conclusions based on a comparison of two species or from the scarce existing literature, the results presented here clearly indicate that bird species living in the same area and using the same habitats and nest sites may still harbor a significantly different bacterial load on their feathers. These findings are therefore consistent with the conclusion of Goodenough and Stallwood (2010) that despite substantial intraspecific variation in bacterial microflora in birds, there are significant interspecific differences even when host species are closely related, ecologically similar, sympatric, and construct very similar nests. Hence, plumage bacterial communities are not solely determined by the soil and habitat characteristics; certain species-specific factors are also important.

4.2. The effect of body topography

A significant correlation was found between the densities of free-living bacteria on the ventral and dorsal feathers of individual Great Tits (III). This matches the finding of Gunderson et al. (2009) who found a strong correlation between bacterial intensities on different body parts of the same Eastern Bluebird individuals. Such results are unsurprising, given that bacterial loads are known to be correlated even between the different parents within breeding pairs (Lucas et al. 2005, Gunderson et al. 2009, I). The most plausible explanation for this phenomenon is that pair-mates infect each other with bacteria via their common nest and brood (I). Hence, if bacteria can easily pass between frequently interacting individuals, it seems reasonable to expect that they can easily spread from one body region to another. It is noteworthy that the correlation between the densities of attached bacteria in different body regions was weaker than the relationship between free-living bacterial densities and was not significant (III). This corresponds to the finding (I) that correlation in the levels of contamination between members of a breeding pair is also much weaker for attached than free-living bacteria. Thus it seems that attached bacteria do not have the same propensity to spread as free-living bacteria.

A rather unexpected result of the study (III) was that densities of feather bacteria associated with Great Tits were higher on the dorsal than the ventral plumage. This is at odds with the prediction made in the Introduction. Significant body topographic variation in feather bacterial loads in birds has been also detected by earlier studies (Burt & Ichida 1999, Bisson et al. 2007). However, the only study (Burt & Ichida 1999) that examined the direction of this difference in several bird species (but not in Great Tits) reported that bacterial loads tend to be higher on ventral than dorsal feathers. Burt and Ichida's (1999) result is somewhat more intuitive than that reported here from Great Tits, as the main source of bacteria is presumed to be the soil, and ventral

feathers certainly come into close contact with the ground and other contaminated substrates. Moreover, the fact that sunlight inhibits bacterial growth (Saranathan & Burt 2007) should also decrease dorsal bacterial densities. However, the yellow chest in Great Tits is probably used to signal to conspecifics (Hörak et al. 2001), and dirt accumulation can reduce plumage coloration (Surmacki & Nowakowski 2007). It has been shown that ornamented bird species devote more time to sanitation behaviours compared with non-ornamented species (Walther & Clayton 2005). Consequently, it is possible that Great Tits preen their chests more carefully than their backs. It is also relevant to note that Burt and Ichida (1999) only examined the presence or absence of feather-degrading bacilli (mainly *Bacillus licheniformis*), which represent only a fraction of the entire diversity of feather bacteria. In contrast, non-selective methods were used in study III to estimate the total abundance of all types of feather-inhabiting bacteria. Thus, it is possible that the difference between ventral and dorsal body parts described by Burt and Ichida (1999) may not hold universally for all ecological types of bacteria, and that other factors besides contact with the ground may be responsible for body topographic variation in the abundance of feather bacteria. The results described here also deliver a methodological message of great importance: in order to ensure comparability, the collection of feather samples must be standardized; otherwise the uneven distribution of bacteria in plumage could generate highly biased results.

4.3. Habitat differences

Habitat-related differences in bacterial density (I, IV) and phylotypic richness (I) in Great Tit plumage showed contrasting patterns: while the number of phylotypes per bird was higher in coniferous habitat, bacterial densities were higher in deciduous habitat. Hence, the habitat-related pattern of variation in feather bacterial densities conforms to the prediction made in the Introduction that birds inhabiting more diverse habitats harbor more bacteria on their feathers. However, habitat-related differences in the phylotypic richness of bacteria did not conform to this pattern.

A negative relationship between species diversity and abundance variables has previously been described for plant communities at relatively high productivity levels (Adams 2009). In bacterial communities, such a relationship can most plausibly be explained by interspecific antagonism and dominance. For example, many bacterial species produce antibacterial chemicals that suppress the growth of other bacteria (Riley & Wertz 2002, Peralta-Sanchez et al. 2010). When fast-breeding (or fast-growing) generalist species (dominants) are present in a community, they might depress overall bacterial diversity (Martin-Platero et al. 2006, Soler et al. 2008, Peralta-Sanchez et al. 2010). Although deciduous forests are generally much more diverse habitats than managed conifers and contain more diverse microhabitats in which birds might

become contaminated with bacteria, rapid infestation with dominant bacteria may inhibit colonization by other bacterial species.

The results of the study (I) indicate that the structure of bacterial communities may vary significantly between habitats even at small geographical scales. In this context it is noteworthy that the physiological condition of breeding Great Tits in the same study area was found to be worse in deciduous than in coniferous forest (Kilgas et al. 2006, 2007, Mägi et al. 2009). Hence, the possibility cannot be ruled out that the relatively high bacterial load associated with deciduous habitat may represent one of the factors contributing to the habitat difference in adult condition.

No significant effect of habitat type on plumage bacterial community was detected in the Pied Flycatcher (II). However, this may be a result of the significantly lower sample sizes collected for this species, compared with Great Tits (II).

4.4. Seasonal changes

One prediction in the Introduction was that the warm and humid environment of the nest and a reduced preening effort due to the increased need to devote more time to feeding offspring would cause a seasonal increase in bacterial abundance on feathers. This prediction was not supported by the results of the study. In fact, female Great Tits supported significantly more bacteria (both attached and free-living) in their plumage during the nest-building period than during the first and second broods (I).

Such a pre-laying peak in bacterial density may be caused by different mechanisms. First, it may be related to nest-building behavior, which brings birds into increased contact with the ground and nest materials (as suggested in I), while possibly leaving them little time for self-preening. Indeed, Great Tits spend considerable time on the ground during the nest-building phase to collect moss, dry grass, hair, wool etc. (Cramp & Perrins 1993). This explanation assumes that the density of bacteria in plumage increases rapidly during the short period of nest building. However, an alternative explanation is derived from the fact that Great Tits roost in cavities (nest boxes in our study area) during the winter. It is therefore possible that damp nest-boxes containing old nest material harbor a diverse bacterial community and leave tits highly infested in the spring (I).

The results of the study (IV) provide support to the first explanation. The pre-laying peak in bacterial density appears to be related to nest-building behavior: the densities of attached feather bacteria in female Great Tits increased during a fairly short period between nest initiation and nest completion. A similar trend, though not statistically significant, was found in the case of free-living bacteria. Both of these trends appeared to be very similar within both habitat types. It is noteworthy that the number of free-living

bacteria tended to increase when the nests were only ‘half ready’ – just after the first animal hairs appeared in the nests. However, the number of attached bacteria tended to increase once nests were completed (IV). This appears to be a logical result, as attached bacteria presumably take some time to become attached.

Thus these results indicate that the high pre-laying density of bacteria in the plumages of Great Tits is not a relic from an earlier period, but appears to be related to nest-building activity. Besides the reasons already listed above, it is perhaps noteworthy that the start of egg-laying in tits is often induced by a sudden spell of warm weather (pers. obs.), and birds may need to hurry with nest-building to such a degree that they have to significantly reduce self-preening during that time. Thus, increase of plumage bacterial load during the short period of nest-building may be one mechanism mediating the costs of nest-building on individuals. Recent studies have shown that nest building in Great Tits and Blue Tits is indeed an energy-demanding endeavor, the outcome of which may be related to individual quality (Tomás et al. 2006, Broggi & Senar 2009). Bacterial abundance in the plumage may decline in subsequent breeding stages due to regular preening activities and reduced contact with the soil.

These results indicate that the density of bacteria in plumage is variable and can fluctuate rapidly during the breeding season. This is also the first study to demonstrate that bacterial load in plumage increases significantly throughout the nest-building process in a free-living arboreal bird. Previously, fast changes in bacterial densities within a single breeding phase have been reported from chick-rearing European Starlings, where free-living, but not attached, bacterial densities increased in response to brood-size manipulation (Lucas et al. 2005).

However, more consistently with the initial prediction, there was also an increase in attached bacterial load between the first and second breeding attempts in the Great Tit (I). This can most plausibly be explained by the extended exposure of plumage to various kinds of microbes, while favorable climatic conditions prevailing in midsummer during the second broods might also enhance bacterial growth and density. Moreover, bacterial density may increase during the season as a result of the cumulative negative effects of reproductive effort on individual condition and preening activities during multiple breeding attempts.

4.5. Annual variation

The density of attached bacteria was significantly higher in 2008 than in 2007 (I). The most plausible reason for the difference between years was the considerably higher mean ambient temperature and level of precipitation in the early spring of 2008 (according to the Estonian Meteorological and Hydrological Institute), which probably favored bacterial growth (Burt & Ichida 1999, 2004, Peele et al. 2009). Hence, the results of the study (III) support the predictions made in the Introduction.

A strong correlation was found between attached bacterial densities within the same individual females in two successive years (III). These results indicate that knowing the bacterial density on feathers of an individual in one particular year does not allow the absolute bacterial density on its feathers in the next year to be estimated. Rather, it allows the rank or magnitude of bacterial load compared with other individuals to be predicted, because this appears to remain fairly constant between years. Hence, bacterial loads of individuals are not merely the result of unpredictable contingencies, but reflect something about individual birds that remains constant over time. Unfortunately, owing to the small sample sizes, these data do not allow us to determine if this is related to individual differences in body condition (Clayton 1999), preening behavior (Walther & Clayton 2005), uropygial oil production or its composition (Martin-Vivaldi et al. 2009, Møller et al. 2009), properties of nesting areas, or to some other factor.

4.6. Sex differences

Comparing the different Great Tit parents within individual breeding pairs showed that bacterial density was correlated within-pair, but females had a tendency on average to support fewer bacterial phylotypes than males (I). As within-pair differences in parental physical activity are expected to be smaller than differences between pairs (Verhulst & Tinbergen 1997), and there is also an expected trade-off between parental provisioning effort and self-preening behavior, one would expect within-pair bacterial densities to be correlated (see also Lucas et al. 2005). An alternative explanation could be that breeding partners share the same breeding territory, are thus exposed to similar bacterial assemblages and may even infect each other with bacteria via their common nest and brood (I). Nonetheless, females of both studied species carried more attached bacteria on their feathers than males (II). In Pied Flycatchers, but not in Great Tits, females also carried more bacterial phylotypes per individual than males (II). Similarly, a higher bacterial load in females, compared with males, was found in European Starlings (Lucas et al. 2005) and Barn Swallows *Hirundo rustica* (Møller et al. 2009, Czirjak et al. 2010). This sex difference is again predictable, because females carry out more diverse activities during the different stages of reproduction (see Introduction). Furthermore, female Great

Tits and Pied Flycatchers in this study area roost in nest boxes, while males roost in the tree canopy. Therefore, females may come into more frequent contact with bacteria-contaminated nest material than males do. Secondly, given that on average male parents invest less effort in their broods than females do (Cramp & Perrins 1993, Verhulst & Tinbergen 1997, Sanz et al. 2000, Sisask et al. 2010), they can devote more time to self-preening.

However, the reasons for significant differences between partners in bacterial phylotypic richness are less clear. Unlike Pied Flycatchers, the number of bacterial phylotypes in the plumage of Great Tits was higher in males than in females (**I**). It seems counterintuitive that male Great Tits support more bacterial phylotypes than females (**I**), for the reasons listed above. However, it may be that because female Great Tits come into greater contact with soil and bacteria than males do, the dominant bacterial species suppress bacterial diversity in their plumage to a greater extent. While males can devote more time to self-preening than females, the application of preen waxes may inhibit the growth and density of some bacteria on feathers, but may not greatly reduce the species diversity of the microbial assemblages (Shawkey et al. 2003a). It is also noteworthy that the difference between the sexes in the mean phylotypic richness of bacteria was larger during second broods than during first broods (**I**). This was presumably related either to the higher ambient temperature during the second broods promoting bacterial development, or to the increased time available for preening in males due to the relatively small size of second broods. The latter is also supported by the finding that males indeed invest relatively less into smaller broods (Verhulst & Tinbergen 1997). However, it then remains unclear why a similar sex \times breeding stage interaction was not revealed in bacterial density.

Together with the few results available from other studies, the results described here suggest that the abundance of feather bacteria in different passerines tends to be generally higher in females than in males. However, as repeatedly shown in this thesis, a greater density of bacteria on feathers does not necessarily also mean a higher number of bacterial phylotypes.

4.7. Bacteria, female condition and breeding parameters

As expected, a significant negative correlation was found between the phylotypic richness of plumage bacteria and adult body mass in Pied Flycatchers (**II**). There was also a negative correlation between attached bacterial density and female mass during the first broods in Great Tits (**I**). However, free-living bacterial density in the latter species was positively correlated with female mass (**I**). The first two findings are thus consistent with the assumption of less contaminated birds also being high-quality individuals that exhibit good body condition (see Introduction). Several studies have

reported similar negative associations between parameters of plumage bacterial assemblages and adult body condition in other species. For example, Gunderson et al. (2009) reported a negative correlation between plumage bacteria intensity and body condition of females in Eastern Bluebirds.

However, there is some inconsistency between the results of different correlative studies. For example, in the same study by Gunderson and colleagues (2009), the body condition of males was in fact positively correlated with bacteria intensity. The counterintuitive positive correlation between free-living bacteria and female mass in Great Tits (**I**) may indicate that female breeders with high body mass (which are usually high-quality individuals; e.g., Merilä & Wiggins 1997) spend more time and energy searching for calcium-rich snail shells for their eggs and food for their nestlings (Tinbergen & Dietz 1994), compared with poor-quality individuals, and are therefore more frequently in contact with possible sources of micro-organisms. Thus, although one certainly cannot draw general conclusions from a small number of correlative studies, the available evidence suggests that the negative relationship between bacteria and body condition may not be universal, and its precise manifestation may depend on various factors, including species-specific factors.

In Great Tits, a negative correlation also emerged between the density of attached bacteria in the plumage of parents and the number of fledged young (**II**). There was no such relationship with clutch size, suggesting that the negative relationship reflects processes occurring during the incubation and/or fledging stage. It may be that birds with high bacterial load are low-quality individuals that did not adjust their brood size to reflect their own reproductive potential and the availability of resources (e.g. Slagsvold & Lifjeld 1990); and such individuals are therefore unable to care adequately for all of their offspring. Indeed, the finding that attached bacterial density and female mass are negatively correlated in Great Tits (**I**) and other species (see discussion and references above) are all in good correspondence with this hypothesis. Secondly, the parents of the broods with high nestling mortality may have increased their parental provisioning effort in order to avoid starvation of their young, and therefore had less time for self-preening. Indeed, it has been previously shown in the same study area, that poor growth of Great Tit nestlings is associated with higher feeding frequency by parents (Mägi et al. 2009). Moreover, Lucas et al. (2005) demonstrated the existence of a trade-off between parental effort and self maintenance experimentally: European Starlings with enlarged broods had more free-living bacteria on their feathers than birds with reduced broods. A third potential explanation for the negative relationship between bacterial load in adults and the number of fledglings is that mortality of nestlings causes an increase in the number of bacteria in the nest (e.g., via the decay of corpses in the nest), and that adult birds that are in contact with such a nest accumulate higher levels of contamination. However, this explanation has to be considered unlikely, because no association between bacterial loads of adults and mortality of nestlings was detected up to the time of adult sampling (**I**).

4.8. Bacteria and plumage coloration

Feather chroma of female Great Tits tended to be negatively related to the phylotypic richness of feather-degrading bacteria during the chick-rearing period, but not during the period before egg-laying (V). At the same time, no associations between chroma and bacterial densities were found in either of the breeding stages. However, the seasonal change in attached bacterial densities was negatively correlated with seasonal change in chroma (V), implying that the increase in bacterial load was accompanied by a decrease in chroma.

To my knowledge, this is the first study on female birds reporting associations between carotenoid-based coloration and plumage bacteria. The only previous study found that male House Finches with redder plumage had lower feather-degrading bacterial loads (Shawkey et al. 2009a). The chroma of the yellow chest plumage of female Great Tits may signal some aspect of individual quality (Senar et al. 2008; Broggi and Senar 2009). Previous studies have found that plumage chroma is more sensitive to developmental perturbations and body condition than hue (Shawkey et al. 2003b, Senar et al. 2008). Chroma in Great Tits has been found to be unrelated to the carotenoid content of the feather, indicating that other mechanisms may be responsible for the variation in this trait (Senar et al. 2008).

Due to the correlative nature of the study (V) it remains unclear whether plumage bacteria directly affect plumage color or if these two variables are correlated with each other through some other factor. For example, it has previously been found that in Eastern Bluebirds feather degrading bacteria can directly affect structural plumage coloration (Shawkey et al. 2007, Gunderson et al. 2009). It is thus possible that feather-degrading bacteria affect the feather microstructure that is also involved in generating the carotenoid-based coloration (Shawkey & Hill 2005). It is also possible that feather-degrading bacteria could damage the carotenoid structure (as suggested by McGraw & Hill 2004).

Alternatively, it has been suggested that moulting may decrease the bacterial load on feathers (Burt & Ichida 1999, but see Giraudeau et al. 2010). At the same time the chroma of new feathers may differ from old feathers reflecting nutritional conditions during the molt (e.g., Hill 2000). As Great Tits start moulting at the end of the breeding season, this process can sometimes overlap with the timing of late broods (e.g., Orell & Ojanen 1980). In the year when plumage coloration was analysed, females were captured before egg-laying, resulting in the desertion of initial nests (V) and presumably also in some delay to egg-laying in replacement nests. Therefore, the possibility cannot be excluded that the negative association between seasonal change in the density of attached bacteria and plumage color was caused by the fact that some individuals started moulting earlier than others and this affected both traits simultaneously.

In several species plumage coloration can also change between molts due to bleaching, dirt accumulation and abrasion. Such changes can also vary individually, so that in some birds plumage coloration increases, while in others it decreases or does not change with time (McGraw & Hill 2004). It is possible that some individuals are able to take better care of their plumage, thus affecting both plumage coloration and bacterial communities living on feathers. In conclusion, whatever the reasons may be, the results of this study showed that the yellow chest color in Great Tits is related to feather-degrading bacterial phylotypic richness and that there are parallel seasonal changes in plumage color and bacterial load.

4.9. Conclusions

This thesis revealed that bacterial assemblages on the feathers of breeding birds are correlated with many avian life-history traits: (i) bacterial load in deciduous habitat (which in this study area appears to be less suitable for breeding than coniferous habitat judging from the physiological condition of adults and fledglings; Kilgas et al. 2006, 2007, Mägi et al. 2009) is higher than that in coniferous habitat (**I**); (ii) females (which carry out a greater diversity of activities during the breeding season) of both studied species carry more bacteria on their feathers than males (**I**, **II**); (iii) bacterial abundance and assemblage richness are correlated with bird body mass (**I**, **II**) and feather color intensity (**V**); and last but not least, (iv) a rapid increase in bacterial abundance occurs during the nest-building period (**IV**). Among other things, these findings suggest that a correlation is present between breeding effort and self-maintenance. Because of the correlative nature of studies in this thesis, one cannot ascertain whether this is a causal trade-off or just correlation, caused or mediated by other factors. However, taking into account earlier findings reported in literature – some of which were based on experimental manipulation (Lucas et al. 2005) (Table 1), or provided indirect evidence that feather-degrading bacteria might actively degrade feathers on living birds (Shawkey et al. 2007, Gunderson et al. 2009, Shawkey et al. 2009a) – the occurrence of such trade-offs seems likely.

Nonetheless, the generality of these findings remains unclear. Between-species and body topographic comparisons showed that certain species-specific factors play an important role, and that, consequently, formerly described patterns are unlikely to apply universally for all bird species and ecological types of bacteria (**II**, **III**). Further study is needed, involving more bird species from different regions and habitats, the use of manipulative approaches, the consideration of a wider range of variables, larger sample sizes, and consideration of bacterial species composition in samples.

SUMMARY

Microorganisms have been shown to play an important role in shaping the life-histories of animals. It has recently been suggested that feather-degrading bacteria influence the trade-off between parental effort and self-preening behavior in birds and may thus affect the individual fitness of birds. However, in order to design appropriate experiments to test this assumption, it is first necessary to collect more information on the basic parameters of these bacterial communities, such as patterns of natural variation in their density and composition, and environmental sources of infestation for birds.

In this thesis, a complex study design was used, coupling molecular and microbiological techniques with field-based collection of ecological and life-history data on two cavity-breeding passerines. The study was conducted in wild breeding populations of Great Tits (*Parus major*) and Pied Flycatchers (*Ficedula hypoleuca*) in a unique study area consisting of a mosaic of two very different habitat types in Estonia, northern Europe. While the species are similar in terms of their nest site requirements, they differ in certain other traits. Environmental and life-history correlates of natural variation in the abundance of free-living and attached bacteria and in the species diversity of feather-degrading bacteria inhabiting bird plumage were explored. Associations of bacterial infestation of feathers with bird body topography, plumage coloration, individual condition and reproductive output was also investigated. The density and species richness of bacterial assemblages was measured using flow cytometry and ribosomal intergenic spacer analysis (RISA).

The density of plumage bacteria was significantly higher in Great Tits than in Pied Flycatchers, providing evidence that the level of bacterial contamination differs even between co-occurring host species that share habitat, nest site and general foraging preferences (paper II).

The densities of both types of bacteria were higher on the dorsal than on the ventral feathers of Great Tits. The densities of free-living, but not attached, bacteria on the two body regions were highly positively correlated (paper III). This result highlights the importance of following a standard sample collection methodology in order to ensure comparability of data.

In Great Tits, the number of bacterial phylotypes per bird was higher in coniferous habitat, while bacterial densities were higher in deciduous habitat (paper I). This demonstrates that the structure of bacterial communities may vary significantly between habitats even at small geographical scales.

This is the first study to demonstrate that the density of attached bacteria on feathers may increase rapidly during the nest building process, and thereafter decline (papers I and IV). The density of free-living bacteria exhibited a similar pattern. The density of bacteria in bird plumage varied significantly between years; however, attached bacterial densities on the same individuals in successive years were correlated (paper III).

Bacterial species richness in Great Tits was sex dependent, with more diverse bacterial assemblages present on males than on females (paper I). At the same time, bacterial densities were higher in females than in males in both species (papers I and II). The latter finding supports the hypothesis that bacterial abundance on the plumage of the different sexes is related to the different activities carried out by males and females during the breeding season.

In Great Tits, free-living bacterial density was positively correlated with female mass; conversely, there was a negative correlation between attached bacterial density and female mass during the period of peak reproductive effort (paper I). In Pied Flycatchers, a negative correlation between parental body mass and the richness of feather-degrading bacterial phylotypes was found (paper II). In Great Tits, higher densities of bacteria in the plumage of parent birds were also associated with the production of fewer fledglings (paper II). These results indicate that feather-bacterial assemblages may be related to bird condition and reproductive effort.

During the chick-rearing period, feather chroma was negatively related to feather-degrading bacterial species richness in Great Tit females. Moreover, the seasonal change in the density of attached bacteria was accompanied by an opposite change in feather chroma (paper V). This is the first study to demonstrate the existence of associations between carotenoid-based coloration and bacterial assemblages in the plumage of female birds.

In conclusion, the findings contained in this thesis will supposedly improve our understanding of how bacterial assemblages on the feathers of breeding birds interact with environmental variables and host life history parameters. Such knowledge is crucial for future attempts to describe and understand the causality of such relationships.

SUMMARY IN ESTONIAN

Sulestiku bakterikoosluste pesitsusaegne varieeruvus kahel vabaltelaval värvuliseliigil

Mikroorganismid mängivad loomade elukäigu kujundamisel tähtsat rolli. Hiljuti on väidetud, et sulgilagundavad bakterid põhjustavad lõivsuhet lindude sigimispingutuse ja sulestiku eest hoolitsemise vahel ja võivad seega avaldada mõju lindude kohasusele. Selleks aga, et osata planeerida korrektsed eksperimente nimetatud väite kontrollimiseks, tuleb esmalt koguda rohkem informatsiooni sulestiku bakterikoosluste põhiparameetrite – arvukuse ja liigilise koosseisu – loodusliku varieeruvuse seaduspärasuste ning lindude bakteritega nakatumise peamiste allikate kohta.

Käesolevas töös kasutati kompleksset uurimistöö ülesehitust, ühendades omavahel molekulaarsete ja mikrobioloogiliste meetodite rakendamise ja ulatusliku ökoloogilise ja elukäigu-andmestiku kogumise kahe suluspesitseva värvulise – rasvatihase (*Parus major*) ja must-kärbsenäpi (*Ficedula hypoleuca*) – looduslikest populatsioonidest, mis paiknevad ühisel, kahe väga erineva elupaiga mosaiiki sisaldaval uurimisalal. Neid liike ühendavad sarnased pesapaigavaliku eelistused, kuid nad erinevad teineteisest mitmete muude omaduste poolest. Uuriti lindude sulgedel elavate vabaltelavate ja kinnitunud bakterite arvukuse ja liigilise mitmekesisuse seoseid elukeskkonna omaduste ja elukäigu parameetritega. Samuti pöörati tähelepanu sulebakterite koosluste omaduste seostele linnu kehapiirkonna, sulestiku värvuse, isendi konditsiooni ja sigimisedukusega. Bakterite asustustiheduse ja bakterikoosluste liigilise mitmekesisuse määramiseks kasutati vastavalt läbivoolutsütomeetriat ja ribosomaalse geenidevahelise speisserjärjestuse analüüsi (RISA).

Sulestikubakterite arvukus leiti rasvatihasel olevat tunduvalt kõrgem kui must-kärbsenäpil. See näitab, et sulestiku bakteritega asustatuse tihedus võib erineda isegi koosesinevate, samu elupaiku, pesapaiku ja toitumisharjumusi jagavate linnuliikide vahel (artikkel II).

Mõlema eelmainitud bakteritüübi arvukus oli rasvatihase seljapoolel kõrgem kui kõhupoolel. Vabaltelavate (kuid mitte kinnitunud) bakterite arvukuses oli kahe kehapiirkonna vahel tugev positiivne seos (artikkel III). See leid rõhutab proovide kogumise meetodika standardiseerimise tähtsust.

Rasvatihastel leiti keskmine sulebakterite mitmekesisus isendi kohta olevat suhteliselt kõrgem okasmetsas, kuid bakterite arvukus oli kõrgem hoopiski lehtmetsas. See tulemus näitab, et sulestikubakterite koosluste omadused võivad erineda elupaikade vahel isegi väga väikesel geograafilisel skaalal.

Käesolevas uurimuses näidati esimest korda, et kinnitunud bakterite arvukus lindude sulestikus võib pesa ehitamise jooksul kiirelt tõusta ja hiljem uuesti langeda (artiklid I ja IV). Sarnased muutused leidsid aset ka vabaltelavate bakterite arvukuses. Bakterite arvukus sulestikus varieerus märkimisväärselt ka

aastate vahel, ehkki kinnitunud bakterite arvukused samadel isenditel eri aastatel olid omavahel korrelatsioonis (artikkel III).

Bakterite liigiline mitmekesisus rasvatihaste sulgedel oli soost sõltuv, kusjuures isastel olid bakterikooslused keskmiselt mitmekesisemad kui emastel lindudel (artikkel I). Samas aga oli bakterite arvukus kõrgem emaste sulgedel, kusjuures see seos kehtis nii rasvatihasel kui ka must-kärbsenäpil (artiklid I ja II). Neist leidudest viimane on kooskõlas hüpoteesiga, mille kohaselt bakterite arvukus eri soost lindude sulgedel on seotud sugupoolte erineva tööjaotuse ja sellest tingitud käitumisega sigimisperioodil.

Vabaltelavate bakterite arvukus rasvatihaste sulgedel oli positiivses seoses emaslinnu massiga, samas aga kinnitunud bakterite arvukus seostus sigimispingutuse tipp-perioodil emase massiga negatiivselt (artikkel I). Must-kärbsenäpidel leiti negatiivne seos linnu kehamassi ja sulebakterite liigilise mitmekesisuse vahel (artikkel II). Rasvatihastel oli bakterite arvukus kõrgem ka neil lindudel, kellel lennuvõimestus vähem järglasi (artikkel II). Need tulemused viitavad, et sulestiku bakterikooslused võivad sõltuvalt lindude konditsioonist ja sigimispingutuse suurusest.

Poegade kasvatamise perioodil oli emaste rasvatihaste sulestiku eredus negatiivses seoses sulgi lagundavate bakterite liigilise mitmekesisusega. Veelgi enam, kinnitunud bakterite arvukuse muutusega pesitsushooaja jooksul kaasnes vastupidise suunaga muutus sulestiku ereduses (artikkel V). Tegu on esimese uurimusega, kus karotinoididel põhineva sulestikuvärvuse seos sulestiku bakteriflooraga leidis kinnitust emaslindude puhul.

Kokkuvõttes aitavad käesolevas töös kirjeldatud tulemused paremini mõista, kuidas sulestikku asustavad bakterikooslused on seotud elukeskonna omadustega ja lindude elukäigu parameetritega. Need teadmised on hädavajalikud edasiste uuringute planeerimiseks, et välja selgitada ja mõista kirjeldatud seoste taga peituvaid põhjuslikke mehhanisme.

REFERENCES

- Adams J. 2009. Species richness: Patterns in the diversity of life., Springer-Praxis, Berlin-Heidelberg-New York.
- Alatalo R. V., Alatalo R. H. 1979. Resource partitioning among a flycatcher guild in Finland. *Oikos* 33: 46–54.
- Bisson I. A., Marra P. P., Burt Jr E. H., Sikaroodi M., Gillevet P. M. 2007. A molecular comparison of plumage and soil bacteria across biogeographic, ecological, and taxonomic scales. *Microbial Ecology* 54: 65–81.
- Bisson I. A., Marra P. P., Burt Jr E. H., Sikaroodi M., Gillevet P. M. 2009. Variation in plumage microbiota depends on season and migration. *Microbial Ecology* 58: 212–220.
- Broggi J., Senar J. C. (2009) Brighter Great Tit parents build bigger nests. *Ibis* 151: 588–591
- Bruce J., Drysdale E. M. 1994. Trans-shell transmission. In: Board R.G., Fuller R. (eds.) *Microbiology of Avian Eggs*. pp. 63–91. Chapman & Hall, London.
- Brush A. 1965. Energetics, temperature regulation and circulation in resting, active and defeathered californian quail, *Lophortyx californicus*. *Comparative Biochemistry and Physiology - A Molecular and Integrative Physiology* 15: 399–421.
- Burt E. H., Ichida J. M. 1999. Occurrence of feather-degrading bacilli in the plumage of birds. *Auk* 116: 364–372.
- Burt E. H., Ichida J. M. 2004. Gloger's rule, feather-degrading bacteria, and color variation among song sparrows. *Condor* 106: 681–686.
- Burt E. H. 2009. A future with feather-degrading bacteria. *J. Avian Biol.* 40: 349–351.
- Burt E. H., Schroeder M. R., Smith L. A., Sroka J. E., McGraw K. J. 2011. Colourful parrot feathers resist bacterial degradation. *Biology Letters* 7: 214–216
- Clayton D. H., Moore J. 1997. Host–parasite evolution: General principles and avian models. Oxford University Press., Oxford.
- Clayton D. H. 1999. Feather-busting bacteria. *Auk* 116: 302–304.
- Cramp S., Perrins C. M. 1993. *The birds of the western palearctic*. Oxford University Press, Oxford-New York.
- Cristol D. A., Armstrong J. L., Whitaker J. M., Forsyth M. H. 2005. Feather-degrading bacteria do not affect feathers on captive birds. *Auk* 122: 222–230.
- Czirják G. A., Anders A. P., Mousseau T. A., Heeb P. 2010. Microorganisms Associated with Feathers of Barn Swallows in Radioactively Contaminated Areas Around Chernobyl *Microbial Ecology* 60:373–380
- Giraudeau M., Gzirják G.A., Duval C., Guiterrez C., Bretagnolle V., Heeb P. 2010. No detected effect of moult on feather bacterial loads in mallards *Anas platyrhynchos*. *Journal of Avian Biology* 41: 678–680.
- Goldstein G., Flory K. R., Browne B. A., Majid S., Ichida J. M., Burt Jr. E. H. 2004. Bacterial degradation of black and white feathers. *The Auk* 121: 656–659.
- Goodenough A. E., Stallwood B. 2010. Intraspecific variation and interspecific differences in the bacterial and fungal assemblages of blue Tit (*Cyanistes caeruleus*) and Great Tit (*Parus major*) nests. *Microbial Ecology* 59: 221–232.
- Gosler A. 1993. *The Great Tit*. Hamlyn Limited, London.
- Grande J. M., Negro J. J., Torres M. J. 2004. The evolution of bird plumage colouration: a role for feather-degrading bacteria? *Ardeola*, 51: 375–383.

- Gunderson A. R., Frame A. M., Swaddle J. P., Forsyth M. H. 2008. Resistance of melanized feathers to bacterial degradation: Is it really so black and white? *J. Avian Biol.* 39: 539–545.
- Gunderson A. R., Forsyth M. H., Swaddle J. P. 2009. Evidence that plumage bacteria influence feather coloration and body condition of eastern bluebirds *Sialia sialis*. *J. Avian Biol.* 40: 440–447.
- Haartman L. von 1954. Der Trauerfliegenschneider, III. Die Nahrungsbiologie. *Acta Zoologica Fennica* 83: 1–96.
- Hackstein J. H. P., Van Alen T. A. 1996. Fecal methanogens and vertebrate evolution. *Evolution* 50: 559–572.
- Hamilton W. D., Zuk M. 1982. Heritable true fitness and bright birds: A role for parasites? *Science* 218: 384–387.
- Hart B. L. 1997. Behavioural defense. In: Clayton D. H., Moore J., (eds.) *Host-parasite evolution: General principles and avian models*. pp. 59–77. Oxford University Press, Oxford.
- Hill G. E. 2000. Energetic constraints on expression of carotenoid-based plumage coloration. *Journal of Avian Biology* 31: 559–566.
- Hörak P., Mänd R., Ots I., Leivits A. 1995. Egg size in the Great Tit *Parus major*: individual, habitat and geographic differences. *Ornis Fennica* 72, 97–114.
- Hörak P., Ots I., Vellau H., Spottiswoode C., Møller A.P. 2001. Carotenoid-based plumage coloration reflects hemoparasite infection and local survival in breeding Great tits. *Oecologia* 126: 166–173.
- Kilgas P., Tilgar V., Mänd R. 2006. Hematological health state indices predict local survival in a small passerine bird, the Great Tit (*Parus major*). *Physiological and Biochemical Zoology* 79(3): 565–572
- Kilgas P., Tilgar V., Mägi M., Mänd R. 2007. Physiological condition of incubating and brood rearing female great tits *Parus major* in two contrasting habitats. *Acta Ornithologica* 42: 129–136.
- Lessells C. M., Boag P. T. 1987. Unrepeatable repeatabilities: a common mistake. *Auk* 104: 116–121.
- Lucas F. S., Bertru G., Hofle M. G. 2003a. Characterization of free-living and attached bacteria in sediments colonized by *Hediste diversicolor*. *Aquatic Microbial Ecology* 32: 165–174.
- Lucas F. S., Broennimann O., Febbraro I., Heeb P. 2003b. High diversity among feather-degrading bacteria from a dry meadow soil. *Microbial Ecology* 45: 282–290.
- Lucas F. S., Moureau B., Jourdie V., Heeb P. 2005. Brood size modifications affect plumage bacterial assemblages of european starlings. *Mol. Ecol.* 14: 639–646.
- Lundberg A., Alatalo R. V. 1992. *The Pied Flycatcher*. Poyser, London.
- Mägi M., Mänd R. 2004. Habitat differences in allocation of eggs between successive breeding attempts in great tits (*Parus major*). *Ecoscience* 11: 361–369.
- Mägi M., Mänd R., Tamm H., Sisask E., Kilgas P., Tilgar V. 2009. Low reproductive success of great tits in the preferred habitat: A role of food availability. *Ecoscience* 16: 145–157.
- Mahler B., Lopez N. I., Digiacoimo A. G., Reboreda J. C. 2010. Increased plumage darkness of female Shiny Cowbirds *Molothrus bonariensis* in the subtropics: an adaptation to bacterial degradation? *Ibis* 152: 775–781
- Mänd R., Tilgar V., Löhmus A., Leivits A. 2005. Providing nest boxes for hole-nesting birds - does habitat matter? *Biodiversity and Conservation* 14: 1823–1840.

- Martin-Platero A. M., Valdivia E., Ruiz-Rodriguez M., Soler J. J., Martin-Vivaldi M., Maqueda M., Martinez-Bueno M. 2006. Characterization of antimicrobial substances produced by *Enterococcus faecalis* mrr 10–3, isolated from the uropygial gland of the hoopoe (*Upupa epops*). *Appl. Environ. Microbiol.* 72: 4245–4249.
- Martin-Vivaldi M., Ruiz-Rodriguez M., Jose Soler J., Manuel Peralta-Sanchez J., Mendez M., Valdivia E., Manuel Martin-Platero A., Martinez-Bueno M. 2009. Seasonal, sexual and developmental differences in hoopoe *Upupa epops* preen gland morphology and secretions: Evidence for a role of bacteria. *J. Avian Biol.* 40: 191–205.
- Martin-Vivaldi M., Pena A., Peralta-Sanchez J. M., Sanchez L., Ananou S., Ruiz-Rodriguez M., Soler J. J. 2010. Antimicrobial chemicals in hoopoe preen secretions are produced by symbiotic bacteria. *Proceedings of the Royal Society B: Biological Sciences* 277: 123–130.
- McGraw K. J., Hill G. E. 2004. Plumage color as a dynamic trait: carotenoid pigmentation of male House Finches (*Carpodacus mexicanus*) fades during the breeding season. *Canadian Journal of Zoology* 82: 734–738.
- Merilä J., Wiggins D. A. 1997. Mass loss in breeding blue tits: The role of energetic stress. *Journal of Animal Ecology* 66: 452–460.
- Merilä J., Hemborg C. 2000. Fitness and feather wear in the collared flycatcher *Ficedula albicollis*. *J. Avian Biol.* 31: 504–510.
- Møller A. P., Czirjak G. A., Heeb P. 2009. Feather micro-organisms and uropygial antimicrobial defences in a colonial passerine bird. *Functional Ecology* 23: 1097–1102.
- Muyzer G., Dewaal E. C., Uitterlinden A. G. 1993. Profiling of complex microbial-populations by denaturing gradient gel-electrophoresis analysis of polymerase chain reaction-amplified genes-coding for 16s ribosomal-rna. *Appl. Environ. Microbiol.* 59: 695–700.
- Muza M. M., Burt E. H., Ichida J. M. 2000. Distribution of bacteria on feathers of some eastern north american birds. *Wilson Bull.* 112: 432–435.
- Nuttall P. A. 1997. Viruses, bacteria and fungi of birds. In: Clayton D. H., Moore J., (eds.) *Host-parasite evolution: General principles and avian models.* pp. 271–302. Oxford University Press, Oxford.
- Orell M., Ojanen M. 1980. Overlap between breeding and moulting in the Great Tit *Parus major* and Willow Tit *P. montanus* in northern Finland. *Ornis Scandinavica* 11: 43–49.
- Ozawa T., Yamaguchi M. 1986. Fractionation and estimation of particle-attached and unattached *Bradyrhizobium japonicum* strains in soils. *Appl. Environ. Microbiol.* 52: 911–914.
- Peele A. M., Burt Jr E. H., Schroeder M. R., Greenberg R. S. 2009. Dark color of the coastal plain swamp sparrow (*Melospiza georgiana nigrescens*) may be an evolutionary response to occurrence and abundance of salt-tolerant feather-degrading bacilli in its plumage. *Auk* 126: 531–535.
- Peralta-Sanchez J. M., Møller A. P., Martin-Platero A. M., Soler J. J. 2010. Number and colour composition of nest lining feathers predict eggshell bacterial community in barn swallow nests: An experimental study. *Functional Ecology* 24: 426–433.
- Peralta-Sanchez J. M., Møller A. P., Soler J. J. 2011. Colour composition of nest lining feathers affects hatching success of barn swallows, *Hirundo rustica* (Passeriformes: Hirundinidae). *Biological Journal of the Linnean Society* 102: 67–74.
- Perrins C. M. 1979. *British tits.* Collins, London.

- Ranjard L., Poly F., Nazaret S. 2000. Monitoring complex bacterial communities using culture-independent molecular techniques: Application to soil environment. *Research in Microbiology* 151: 167–177.
- Reneerkens J., Versteegh M. A., Schneider A. M., Piersma T., Burt E. H. 2008. Seasonally changing preen-wax composition: Red knots' (*Calidris canutus*) flexible defense against feather-degrading bacteria? *Auk* 125: 285–290.
- Revis H. C., Waller D. A. 2004. Bactericidal and fungicidal activity of ant chemicals on feather parasites: an evaluation of anting behavior as a method of self-medication in songbirds. *The Auk* 121: 1262–1268.
- Riffel A., Lucas F., Heeb P., Brandelli A. 2003. Characterization of a new keratinolytic bacterium that completely degrades native feather keratin. *Archives of Microbiology* 179: 258–265.
- Riley M. A., Wertz J. E. 2002. Bacteriocin diversity: Ecological and evolutionary perspectives. *Biochimie* 84: 357–364.
- Ruiz-de-Castaneda R., Burt E. H., Gonzalez-Braojos S., Moreno J. 2012. Bacterial degradability of an intrafeather unmelanized ornament: a role for feather-degrading bacteria in sexual selection? *Biological Journal of the Linnean Society* 105: 409–419.
- Ruiz-Rodriguez M., Valdivia E., Soler J. J., Martin-Vivaldi M., Martin-Platero A. M., Martinez-Bueno M. 2009. Symbiotic bacteria living in the hoopoe's uropygial gland prevent feather degradation. *Journal of Experimental Biology* 212: 3621–3626.
- Sangali S., Brandelli A. 2000. Isolation and characterization of a novel feather-degrading bacterial strain. *Applied Biochemistry and Biotechnology – Part A Enzyme Engineering and Biotechnology* 87: 17–24.
- Sanz J. J., Kranenbarg S., Tinbergen J. M. 2000. Differential response by males and females to manipulation of partner contribution in the great tit (*Parus major*). *Journal of Animal Ecology* 69: 74–84.
- Saranathan V., Burt E. H. 2007. Sunlight on feathers inhibits feather-degrading bacteria. *Wilson J. Ornithol.* 119: 239–245.
- Selje N., Simon M. 2003. Composition and dynamics of particle-associated and free-living bacterial communities in the weser estuary, Germany. *Aquatic Microbial Ecology* 30: 221–237.
- Senar J. C., Negro J. J., Quesada J., Ruiz I., Garrido J. 2008. Two pieces of information in a single trait? The yellow breast of the great tit (*Parus major*) reflects both pigment acquisition and body condition. *Behaviour* 145: 1195–1210.
- Shawkey M. D., Pillai S. R., Hill G. E. 2003a. Chemical warfare? Effects of uropygial oil on feather-degrading bacteria. *J. Avian Biol.* 34: 345–349.
- Shawkey M. D., Estes A. M., Siefferman L. M., Hill G. E. 2003b. Nanostructure predicts intraspecific variation in ultraviolet-blue plumage colour. *Proceedings of the Royal Society B: Biological Sciences* 270: 1455–1460.
- Shawkey M. D., Hill G. E. 2005. Carotenoids need structural colours to shine. *Biology Letters* 1: 121–124.
- Shawkey M. D., Mills K. L., Dale C., Hill G. E. 2005. Microbial diversity of wild bird feathers revealed through culture-based and culture-independent techniques. *Microbial Ecology* 50: 40–47.
- Shawkey M. D., Hussain M. J., Strong A. L., Hagelin J. C., Vollmer A. C., Hill G. E. 2006. Use of culture-independent methods to compare bacterial assemblages on feathers of crested and least Auklets (*Aethia cristatella* and *Aethia pusilla*) with those of passerines. *Waterbirds*, 29: 507–511.

- Shawkey M. D., Pillai S. R., Hill G. E., Siefferman L. M., Roberts S. R. 2007. Bacteria as an agent for change in structural plumage color: Correlational and experimental evidence. *The American Naturalist* 169: S112-S121.
- Shawkey M. D., Pillai S. R., Hill G. E. 2009a. Do feather-degrading bacteria affect sexually selected plumage color? *Naturwissenschaften* 96: 123–128.
- Shawkey M. D., Firestone M. K., Brodie E. L., Beissinger S. R. 2009b. Avian incubation inhibits growth and diversification of bacterial assemblages on eggs. *PLoS ONE* 4:
- Sisask E., Mänd R., Mägi M., Tilgar V. 2010. Parental provisioning behaviour in pied flycatchers is well adjusted to local conditions in a mosaic of deciduous and coniferous habitat. *Bird Study* 57:447–457.
- Slagsvold T., Lifjeld J. T. 1990. Influence of male and female quality on clutch size in tits (*Parus spp.*). *Ecology* 71: 1258–1266.
- Soler J. J., Martin-Vivaldi M., Ruiz-Rodriguez M., Valdivia E., Martin-Platero A. M., Martinez-Bueno M., Peralta-Sanchez J. M., Mendez M. 2008. Symbiotic association between hoopoes and antibiotic-producing bacteria that live in their uropygial gland. *Functional Ecology* 22: 864–871.
- Stach J. E. M., Maldonado L. A., Masson D. G., Ward A. C., Goodfellow M., Bull A. T. 2003. Statistical approaches for estimating actinobacterial diversity in marine sediments. *Appl. Environ. Microbiol.* 69: 6189–6200.
- Swaddle J. P., Witter M. S., Cuthill I. C., Budden A., McCowen P. 1996. Plumage condition affects flight performance in common starlings: Implications for developmental homeostasis, abrasion and moult. *J. Avian Biol.* 27: 103–111.
- Tinbergen J. M., Dietz M. W. 1994. Parental energy expenditure during brood rearing in the great tit (*Parus major*) in relation to body mass, temperature, food availability and clutch size. *Functional Ecology* 8: 563–572.
- Tomás G., Merino S., Moreno J., Sanz J.J., Morales J., García-Fraile S. (2006) Nest weight and female health in the blue tit (*Cyanistes caeruleus*). *Auk* 123: 1013–1021
- Verhulst S., Tinbergen J. M. 1997. Clutch size and parental effort in the great tit *Parus major*. *Ardea* 85: 111–126.
- Vilbaste H. 1994. Great Tit. In: Leibak E., Lilleht V., Veromann H. (eds.) *Birds of Estonia. Status, distribution and numbers.* Estonian Academy Publishers, Tallinn.
- Walther B. A., Clayton D. H. 2005. Elaborate ornaments are costly to maintain: Evidence for high maintenance handicaps. *Behavioral Ecology* 16: 89–95.

ACKNOWLEDGEMENTS

Several people sacrificed their valuable time to help me complete the work described in this thesis: Veljo Kisand, Natalja Tšertova, Jelena Kiprovskaia and Hannes Luidalepp from the Molecular Microbiology Group of the Institute of Technology, Tartu University taught me new laboratory techniques and helped whenever I needed technical support. The same is true for Urmas Saarma from the Chair of Zoology of our department. Other members of my own working group, Marko Mägi, Priit Kilgas and Elo Rasmann, assisted in field studies. Among them, Priit earns special thanks for his major contributions to papers IV and V. Ants Kaasik gave valuable advice on statistical issues and John Davison did his best to improve the language of the manuscripts. Last but not least (to be honest, most importantly), Raivo Mänd and Vallo Tilgar gave me ideas, contributed to planning studies, helped with methodological issues and statistical analyses and helped to shape my results and ideas into the form of readable manuscripts. In short, I thank them for the nerve-wracking duty of supervising my studies.

Finally, I also want to thank all the other people who have helped me to struggle through my Ph.D. studies, but whose contribution I have shamefully forgotten.

PUBLICATIONS

CURRICULUM VITAE

I. General

Name: Pauli Saag

Date and place of birth: 21.05.1981 Tartu

Citizenship: Estonian

Address: Tartu, Kungla 46, *e-mail*: pauli@ut.ee

Position: University of Tartu, Institute of Ecology and Earth Sciences,
Department of Zoology, Researcer

Educational history

2006 University of Tartu, Faculty of Biology and Geography,
Magister Scientarum in zoology

2004 University of Tartu, Faculty of Biology and Geography, *Baccalaureus* in
zoology

1999 Tartu Mart Reinik High School

Languages spoken: Estonian, English, Russian

Working experience

2010– University of Tartu, Institute of Ecology and Earth Sciences,
Department of Zoology, Researcer

2006–2009 University of Tartu, Institute of Ecology and Earth Sciences,
Department of Zoology, specialist

2003–2004 NGO Eagleclub, expert

II. Research history

1. Research interests: biosystematics ja ecophysiology (ecophysiology,
molecular ecology, molecular systematics and population genetics)

2. Publications:

Kilgas, P., Saag, P., Mägi, M., Edenberg, M., Tilgar, V., Mänd, R. (2012).
Variation in assemblages of feather bacteria in relation to plumage
coloration in female great tits (*Parus major*). *Condor*, xx–xx. [ilmumas]

Saag, P.; Kilgas, P.; Mägi, M.; Tilgar, V.; Mänd, R. (2012). Inter-annual and
body topographic consistency in the plumage bacterial load of Great Tits.
Journal of Field Ornithology, 83(1): 94–100.

- Kilgas, P.; Saag, P.; Mägi, M.; Tilgar, V.; Mänd, R. (2011). Plumage bacterial load increases during nest-building in a passerine bird. *Journal of Ornithology*, doi:10.1007/s10336-011-0801-3 [ilmumas]
- Saag, P., Mänd, R., Tilgar, V., Kilgas, P., Mägi, M., Rasmann, E. (2011). Plumage bacterial load is related to species, sex, biometrics and fledging success in co-occurring cavity-breeding passerines. *Acta Ornithologica*, 46(2), 191–201.
- Tilgar, V.; Moks, K.; Saag, P. (2011). Predator-induced stress changes parental feeding behavior in pied flycatchers. *Behavioral Ecology*, 22(1), 23–28.
- Saag, P.; Tilgar, V.; Mänd, R.; Kilgas, P.; Mägi, M. (2011). Plumage bacterial assemblages in a breeding wild passerine: relationships with ecological factors and body condition. *Microbial Ecology*, 61(4), 740–749.
- Tilgar, V.; Saag, P.; Külavee, R.; Mänd, R. (2010). Behavioural and physiological responses of nestling pied flycatchers to acoustic stress. *Hormones and Behavior*, 57(4–5), 481–487.
- Väli, Ü.; Saag, P.; Dombrowski, V.; Meyburg, B.-U.; Maciorowski, G.; Mizera, T.; Treinys, R.; Fagerberg, S. (2010). Microsatellites and single nucleotide polymorphisms in avian hybrid identification: a comparative case study. *Journal of Avian Biology*, 41(1), 34–49.
- Tilgar, V.; Saag, P.; Moks, K. (2009). Development of stress response in nestling pied flycatchers. *Journal of Comparative Physiology a – Neuroethology Sensory Neural and Behavioral Physiology*, 195(8), 799–803.
- Saag, P.; Paaver, T.; Väli, Ü. (2007). Lack of between- and within-species isoenzyme variability in *Aquila* eagles (Aves: Accipitriformes). *Biochemical Systematics and Ecology*, 36, 774–776.

ELULOOKIRJELDUS

I. Üldandmed

Nimi: Pauli Saag

Sünniaeg ja koht: 21.05.1981 Tartu

Kodakondsus: Eesti

Aadress: Tartu, Kungla 46, *e-mail*: pauli@ut.ee

Praegune töökoht, amet: Tartu Ülikool, Ökoloogia ja maateaduste instituut,
Zooloogia osakond, teadur

Haridus

2006 Tartu Ülikool, Bioloogia-geograafiateaduskond, *Magister Scientarum*
zooloogias

2004 Tartu Ülikool, Bioloogia-geograafiateaduskond, *Baccalaureus* zooloogias

1999 Tartu Mart Reiniku Gümnaasium

Keelteoskus: Eesti, Inglise, Vene

Töökogemus

2010– Tartu Ülikooli loomaökoloogia teadur

2006–2009 Tartu Ülikooli linnuökoloogia spetsialist

2003–2004 MTÜ Kotkaklubi ekspert

II. Teaduslik ja arendustegevus

3. Peamised uurimisvaldkonnad: biosüsteematika ja ökofüsioloogia (ökofüsioloogia, molekulaarne ökoloogia, molekulaarne süsteematika ja populatsioonigeneetika)

4. Publikatsioonide loetelu:

Kilgas, P., Saag, P., Mägi, M., Edenberg, M., Tilgar, V., Mänd, R. (2012). Variation in assemblages of feather bacteria in relation to plumage coloration in female great tits (*Parus major*). *Condor*, xx–xx. [ilmumas]

Saag, P.; Kilgas, P.; Mägi, M.; Tilgar, V.; Mänd, R. (2012). Inter-annual and body topographic consistency in the plumage bacterial load of Great Tits. *Journal of Field Ornithology*, 83(1): 94–100.

Kilgas, P.; Saag, P.; Mägi, M.; Tilgar, V.; Mänd, R. (2011). Plumage bacterial load increases during nest-building in a passerine bird. *Journal of Ornithology*, doi:10.1007/s10336-011-0801-3 [ilmumas]

Saag, P., Mänd, R., Tilgar, V., Kilgas, P., Mägi, M., Rasmann, E. (2011). Plumage bacterial load is related to species, sex, biometrics and fledging

- success in co-occurring cavity-breeding passerines. *Acta Ornithologica*, 46(2), 191–201.
- Tilgar, V.; Moks, K.; Saag, P. (2011). Predator-induced stress changes parental feeding behavior in pied flycatchers. *Behavioral Ecology*, 22(1), 23–28.
- Saag, P.; Tilgar, V.; Mänd, R.; Kilgas, P.; Mägi, M. (2011). Plumage bacterial assemblages in a breeding wild passerine: relationships with ecological factors and body condition. *Microbial Ecology*, 61(4), 740–749.
- Tilgar, V.; Saag, P.; Külavee, R.; Mänd, R. (2010). Behavioural and physiological responses of nestling pied flycatchers to acoustic stress. *Hormones and Behavior*, 57(4-5), 481–487.
- Väli, Ü.; Saag, P.; Dombrowski, V.; Meyburg, B.-U.; Maciorowski, G.; Mizera, T.; Treinys, R.; Fagerberg, S. (2010). Microsatellites and single nucleotide polymorphisms in avian hybrid identification: a comparative case study. *Journal of Avian Biology*, 41(1), 34–49.
- Tilgar, V.; Saag, P.; Moks, K. (2009). Development of stress response in nestling pied flycatchers. *Journal of Comparative Physiology a – Neuroethology Sensory Neural and Behavioral Physiology*, 195(8), 799–803.
- Saag, P.; Paaver, T.; Väli, Ü. (2007). Lack of between- and within-species isoenzyme variability in *Aquila* eagles (Aves: Accipitriformes). *Biochemical Systematics and Ecology*, 36, 774–776.

DISSERTATIONES BIOLOGICAE UNIVERSITATIS TARTUENSIS

1. **Toivo Maimets.** Studies of human oncoprotein p53. Tartu, 1991, 96 p.
2. **Enn K. Seppet.** Thyroid state control over energy metabolism, ion transport and contractile functions in rat heart. Tartu, 1991, 135 p.
3. **Kristjan Zobel.** Epifüütsete makrosamblike väärtus õhu saastuse indikaatoritena Hamar-Dobani boreaalsetes mägimetsades. Tartu, 1992, 131 lk.
4. **Andres Mäe.** Conjugal mobilization of catabolic plasmids by transposable elements in helper plasmids. Tartu, 1992, 91 p.
5. **Maia Kivisaar.** Studies on phenol degradation genes of *Pseudomonas* sp. strain EST 1001. Tartu, 1992, 61 p.
6. **Allan Nurk.** Nucleotide sequences of phenol degradative genes from *Pseudomonas* sp. strain EST 1001 and their transcriptional activation in *Pseudomonas putida*. Tartu, 1992, 72 p.
7. **Ülo Tamm.** The genus *Populus* L. in Estonia: variation of the species biology and introduction. Tartu, 1993, 91 p.
8. **Jaanus Remme.** Studies on the peptidyltransferase centre of the *E.coli* ribosome. Tartu, 1993, 68 p.
9. **Ülo Langel.** Galanin and galanin antagonists. Tartu, 1993, 97 p.
10. **Arvo Käärnd.** The development of an automatic online dynamic fluorescence-based pH-dependent fiber optic penicillin flowthrough biosensor for the control of the benzylpenicillin hydrolysis. Tartu, 1993, 117 p.
11. **Lilian Järvekülg.** Antigenic analysis and development of sensitive immunoassay for potato viruses. Tartu, 1993, 147 p.
12. **Jaak Palumets.** Analysis of phytomass partition in Norway spruce. Tartu, 1993, 47 p.
13. **Arne Sellin.** Variation in hydraulic architecture of *Picea abies* (L.) Karst. trees grown under different environmental conditions. Tartu, 1994, 119 p.
13. **Mati Reeben.** Regulation of light neurofilament gene expression. Tartu, 1994, 108 p.
14. **Urmas Tartes.** Respiration rhythms in insects. Tartu, 1995, 109 p.
15. **Ülo Puurand.** The complete nucleotide sequence and infections *in vitro* transcripts from cloned cDNA of a potato A potyvirus. Tartu, 1995, 96 p.
16. **Peeter Hõrak.** Pathways of selection in avian reproduction: a functional framework and its application in the population study of the great tit (*Parus major*). Tartu, 1995, 118 p.
17. **Erkki Truve.** Studies on specific and broad spectrum virus resistance in transgenic plants. Tartu, 1996, 158 p.
18. **Illar Pata.** Cloning and characterization of human and mouse ribosomal protein S6-encoding genes. Tartu, 1996, 60 p.
19. **Ülo Niinemets.** Importance of structural features of leaves and canopy in determining species shade-tolerance in temperature deciduous woody taxa. Tartu, 1996, 150 p.

20. **Ants Kurg.** Bovine leukemia virus: molecular studies on the packaging region and DNA diagnostics in cattle. Tartu, 1996, 104 p.
21. **Ene Ustav.** E2 as the modulator of the BPV1 DNA replication. Tartu, 1996, 100 p.
22. **Aksel Soosaar.** Role of helix-loop-helix and nuclear hormone receptor transcription factors in neurogenesis. Tartu, 1996, 109 p.
23. **Maido Remm.** Human papillomavirus type 18: replication, transformation and gene expression. Tartu, 1997, 117 p.
24. **Tiiu Kull.** Population dynamics in *Cypripedium calceolus* L. Tartu, 1997, 124 p.
25. **Kalle Olli.** Evolutionary life-strategies of autotrophic planktonic microorganisms in the Baltic Sea. Tartu, 1997, 180 p.
26. **Meelis Pärtel.** Species diversity and community dynamics in calcareous grassland communities in Western Estonia. Tartu, 1997, 124 p.
27. **Malle Leht.** The Genus *Potentilla* L. in Estonia, Latvia and Lithuania: distribution, morphology and taxonomy. Tartu, 1997, 186 p.
28. **Tanel Tenson.** Ribosomes, peptides and antibiotic resistance. Tartu, 1997, 80 p.
29. **Arvo Tuvikene.** Assessment of inland water pollution using biomarker responses in fish *in vivo* and *in vitro*. Tartu, 1997, 160 p.
30. **Urmas Saarma.** Tuning ribosomal elongation cycle by mutagenesis of 23S rRNA. Tartu, 1997, 134 p.
31. **Henn Ojaveer.** Composition and dynamics of fish stocks in the gulf of Riga ecosystem. Tartu, 1997, 138 p.
32. **Lembi Lõugas.** Post-glacial development of vertebrate fauna in Estonian water bodies. Tartu, 1997, 138 p.
33. **Margus Pooga.** Cell penetrating peptide, transportan, and its predecessors, galanin-based chimeric peptides. Tartu, 1998, 110 p.
34. **Andres Saag.** Evolutionary relationships in some cetrarioid genera (Lichenized Ascomycota). Tartu, 1998, 196 p.
35. **Aivar Liiv.** Ribosomal large subunit assembly *in vivo*. Tartu, 1998, 158 p.
36. **Tatjana Oja.** Isoenzyme diversity and phylogenetic affinities among the eurasian annual bromes (*Bromus* L., Poaceae). Tartu, 1998, 92 p.
37. **Mari Moora.** The influence of arbuscular mycorrhizal (AM) symbiosis on the competition and coexistence of calcareous crassland plant species. Tartu, 1998, 78 p.
38. **Olavi Kurina.** Fungus gnats in Estonia (*Diptera: Bolitophilidae, Keroplattidae, Macroceridae, Ditomyiidae, Diadocidiidae, Mycetophilidae*). Tartu, 1998, 200 p.
39. **Andrus Tasa.** Biological leaching of shales: black shale and oil shale. Tartu, 1998, 98 p.
40. **Arnold Kristjuhan.** Studies on transcriptional activator properties of tumor suppressor protein p53. Tartu, 1998, 86 p.

41. **Sulev Ingerpuu.** Characterization of some human myeloid cell surface and nuclear differentiation antigens. Tartu, 1998, 163 p.
42. **Veljo Kisand.** Responses of planktonic bacteria to the abiotic and biotic factors in the shallow lake Võrtsjärv. Tartu, 1998, 118 p.
43. **Kadri Põldmaa.** Studies in the systematics of hypomyces and allied genera (Hypocreales, Ascomycota). Tartu, 1998, 178 p.
44. **Markus Vetemaa.** Reproduction parameters of fish as indicators in environmental monitoring. Tartu, 1998, 117 p.
45. **Heli Talvik.** Prepatent periods and species composition of different *Oesophagostomum* spp. populations in Estonia and Denmark. Tartu, 1998, 104 p.
46. **Katrin Heinsoo.** Cuticular and stomatal antechamber conductance to water vapour diffusion in *Picea abies* (L.) karst. Tartu, 1999, 133 p.
47. **Tarmo Annilo.** Studies on mammalian ribosomal protein S7. Tartu, 1998, 77 p.
48. **Indrek Ots.** Health state indicies of reproducing great tits (*Parus major*): sources of variation and connections with life-history traits. Tartu, 1999, 117 p.
49. **Juan Jose Cantero.** Plant community diversity and habitat relationships in central Argentina grasslands. Tartu, 1999, 161 p.
50. **Rein Kalamees.** Seed bank, seed rain and community regeneration in Estonian calcareous grasslands. Tartu, 1999, 107 p.
51. **Sulev Kõks.** Cholecystokinin (CCK) — induced anxiety in rats: influence of environmental stimuli and involvement of endopioid mechanisms and erotonin. Tartu, 1999, 123 p.
52. **Ebe Sild.** Impact of increasing concentrations of O₃ and CO₂ on wheat, clover and pasture. Tartu, 1999, 123 p.
53. **Ljudmilla Timofejeva.** Electron microscopical analysis of the synaptonemal complex formation in cereals. Tartu, 1999, 99 p.
54. **Andres Valkna.** Interactions of galanin receptor with ligands and G-proteins: studies with synthetic peptides. Tartu, 1999, 103 p.
55. **Taavi Virro.** Life cycles of planktonic rotifers in lake Peipsi. Tartu, 1999, 101 p.
56. **Ana Rebane.** Mammalian ribosomal protein S3a genes and intron-encoded small nucleolar RNAs U73 and U82. Tartu, 1999, 85 p.
57. **Tiina Tamm.** Cocksfoot mottle virus: the genome organisation and translational strategies. Tartu, 2000, 101 p.
58. **Reet Kurg.** Structure-function relationship of the bovine papilloma virus E2 protein. Tartu, 2000, 89 p.
59. **Toomas Kivisild.** The origins of Southern and Western Eurasian populations: an mtDNA study. Tartu, 2000, 121 p.
60. **Niilo Kaldalu.** Studies of the TOL plasmid transcription factor XylS. Tartu 2000. 88 p.

61. **Dina Lepik.** Modulation of viral DNA replication by tumor suppressor protein p53. Tartu 2000. 106 p.
62. **Kai Vellak.** Influence of different factors on the diversity of the bryophyte vegetation in forest and wooded meadow communities. Tartu 2000. 122 p.
63. **Jonne Kotta.** Impact of eutrophication and biological invasions on the structure and functions of benthic macrofauna. Tartu 2000. 160 p.
64. **Georg Martin.** Phytobenthic communities of the Gulf of Riga and the inner sea the West-Estonian archipelago. Tartu, 2000. 139 p.
65. **Silvia Sepp.** Morphological and genetical variation of *Alchemilla L.* in Estonia. Tartu, 2000. 124 p.
66. **Jaan Liira.** On the determinants of structure and diversity in herbaceous plant communities. Tartu, 2000. 96 p.
67. **Priit Zingel.** The role of planktonic ciliates in lake ecosystems. Tartu 2001. 111 p.
68. **Tiit Teder.** Direct and indirect effects in Host-parasitoid interactions: ecological and evolutionary consequences. Tartu 2001. 122 p.
69. **Hannes Kollist.** Leaf apoplastic ascorbate as ozone scavenger and its transport across the plasma membrane. Tartu 2001. 80 p.
70. **Reet Marits.** Role of two-component regulator system PehR-PehS and extracellular protease PrtW in virulence of *Erwinia Carotovora* subsp. *Carotovora*. Tartu 2001. 112 p.
71. **Vallo Tilgar.** Effect of calcium supplementation on reproductive performance of the pied flycatcher *Ficedula hypoleuca* and the great tit *Parus major*, breeding in Northern temperate forests. Tartu, 2002. 126 p.
72. **Rita Hõrak.** Regulation of transposition of transposon Tn4652 in *Pseudomonas putida*. Tartu, 2002. 108 p.
73. **Liina Eek-Piirsoo.** The effect of fertilization, mowing and additional illumination on the structure of a species-rich grassland community. Tartu, 2002. 74 p.
74. **Krõõt Aasamaa.** Shoot hydraulic conductance and stomatal conductance of six temperate deciduous tree species. Tartu, 2002. 110 p.
75. **Nele Ingerpuu.** Bryophyte diversity and vascular plants. Tartu, 2002. 112 p.
76. **Neeme Tõnisson.** Mutation detection by primer extension on oligonucleotide microarrays. Tartu, 2002. 124 p.
77. **Margus Pensa.** Variation in needle retention of Scots pine in relation to leaf morphology, nitrogen conservation and tree age. Tartu, 2003. 110 p.
78. **Asko Lõhmus.** Habitat preferences and quality for birds of prey: from principles to applications. Tartu, 2003. 168 p.
79. **Viljar Jaks.** p53 — a switch in cellular circuit. Tartu, 2003. 160 p.
80. **Jaana Männik.** Characterization and genetic studies of four ATP-binding cassette (ABC) transporters. Tartu, 2003. 140 p.
81. **Marek Sammul.** Competition and coexistence of clonal plants in relation to productivity. Tartu, 2003. 159 p.

82. **Ivar Ilves.** Virus-cell interactions in the replication cycle of bovine papillomavirus type 1. Tartu, 2003. 89 p.
83. **Andres Männik.** Design and characterization of a novel vector system based on the stable replicator of bovine papillomavirus type 1. Tartu, 2003. 109 p.
84. **Ivika Ostonen.** Fine root structure, dynamics and proportion in net primary production of Norway spruce forest ecosystem in relation to site conditions. Tartu, 2003. 158 p.
85. **Gudrun Veldre.** Somatic status of 12–15-year-old Tartu schoolchildren. Tartu, 2003. 199 p.
86. **Ülo Väli.** The greater spotted eagle *Aquila clanga* and the lesser spotted eagle *A. pomarina*: taxonomy, phylogeography and ecology. Tartu, 2004. 159 p.
87. **Aare Abroi.** The determinants for the native activities of the bovine papillomavirus type 1 E2 protein are separable. Tartu, 2004. 135 p.
88. **Tiina Kahre.** Cystic fibrosis in Estonia. Tartu, 2004. 116 p.
89. **Helen Orav-Kotta.** Habitat choice and feeding activity of benthic suspension feeders and mesograzers in the northern Baltic Sea. Tartu, 2004. 117 p.
90. **Maarja Öpik.** Diversity of arbuscular mycorrhizal fungi in the roots of perennial plants and their effect on plant performance. Tartu, 2004. 175 p.
91. **Kadri Tali.** Species structure of *Neotinea ustulata*. Tartu, 2004. 109 p.
92. **Kristiina Tambets.** Towards the understanding of post-glacial spread of human mitochondrial DNA haplogroups in Europe and beyond: a phylogeographic approach. Tartu, 2004. 163 p.
93. **Arvi Jõers.** Regulation of p53-dependent transcription. Tartu, 2004. 103 p.
94. **Lilian Kadaja.** Studies on modulation of the activity of tumor suppressor protein p53. Tartu, 2004. 103 p.
95. **Jaak Truu.** Oil shale industry wastewater: impact on river microbial community and possibilities for bioremediation. Tartu, 2004. 128 p.
96. **Maire Peters.** Natural horizontal transfer of the *pheBA* operon. Tartu, 2004. 105 p.
97. **Ülo Maiväli.** Studies on the structure-function relationship of the bacterial ribosome. Tartu, 2004. 130 p.
98. **Merit Otsus.** Plant community regeneration and species diversity in dry calcareous grasslands. Tartu, 2004. 103 p.
99. **Mikk Heidemaa.** Systematic studies on sawflies of the genera *Dolerus*, *Empria*, and *Caliroa* (Hymenoptera: Tenthredinidae). Tartu, 2004. 167 p.
100. **Ilmar Tõnno.** The impact of nitrogen and phosphorus concentration and N/P ratio on cyanobacterial dominance and N₂ fixation in some Estonian lakes. Tartu, 2004. 111 p.
101. **Lauri Saks.** Immune function, parasites, and carotenoid-based ornaments in greenfinches. Tartu, 2004. 144 p.
102. **Siiri Rootsi.** Human Y-chromosomal variation in European populations. Tartu, 2004. 142 p.

103. **Eve Vedler.** Structure of the 2,4-dichloro-phenoxyacetic acid-degradative plasmid pEST4011. Tartu, 2005. 106 p.
104. **Andres Tover.** Regulation of transcription of the phenol degradation *pheBA* operon in *Pseudomonas putida*. Tartu, 2005. 126 p.
105. **Helen Udras.** Hexose kinases and glucose transport in the yeast *Hansenula polymorpha*. Tartu, 2005. 100 p.
106. **Ave Suija.** Lichens and lichenicolous fungi in Estonia: diversity, distribution patterns, taxonomy. Tartu, 2005. 162 p.
107. **Piret Lõhmus.** Forest lichens and their substrata in Estonia. Tartu, 2005. 162 p.
108. **Inga Lips.** Abiotic factors controlling the cyanobacterial bloom occurrence in the Gulf of Finland. Tartu, 2005. 156 p.
109. **Kaasik, Krista.** Circadian clock genes in mammalian clockwork, metabolism and behaviour. Tartu, 2005. 121 p.
110. **Juhan Javoiš.** The effects of experience on host acceptance in ovipositing moths. Tartu, 2005. 112 p.
111. **Tiina Sedman.** Characterization of the yeast *Saccharomyces cerevisiae* mitochondrial DNA helicase Hmi1. Tartu, 2005. 103 p.
112. **Ruth Aguraiuja.** Hawaiian endemic fern lineage *Diellia* (Aspleniaceae): distribution, population structure and ecology. Tartu, 2005. 112 p.
113. **Riho Teras.** Regulation of transcription from the fusion promoters generated by transposition of Tn4652 into the upstream region of *pheBA* operon in *Pseudomonas putida*. Tartu, 2005. 106 p.
114. **Mait Metspalu.** Through the course of prehistory in india: tracing the mtDNA trail. Tartu, 2005. 138 p.
115. **Elin Lõhmussaar.** The comparative patterns of linkage disequilibrium in European populations and its implication for genetic association studies. Tartu, 2006. 124 p.
116. **Priit Kopper.** Hydraulic and environmental limitations to leaf water relations in trees with respect to canopy position. Tartu, 2006. 126 p.
117. **Heili Ilves.** Stress-induced transposition of Tn4652 in *Pseudomonas Putida*. Tartu, 2006. 120 p.
118. **Silja Kuusk.** Biochemical properties of Hmi1p, a DNA helicase from *Saccharomyces cerevisiae* mitochondria. Tartu, 2006. 126 p.
119. **Kersti Püssa.** Forest edges on medium resolution landsat thematic mapper satellite images. Tartu, 2006. 90 p.
120. **Lea Tummeleht.** Physiological condition and immune function in great tits (*Parus major* L.): Sources of variation and trade-offs in relation to growth. Tartu, 2006. 94 p.
121. **Toomas Esperk.** Larval instar as a key element of insect growth schedules. Tartu, 2006. 186 p.
122. **Harri Valdmann.** Lynx (*Lynx lynx*) and wolf (*Canis lupus*) in the Baltic region: Diets, helminth parasites and genetic variation. Tartu, 2006. 102 p.

123. **Priit Jõers.** Studies of the mitochondrial helicase Hmi1p in *Candida albicans* and *Saccharomyces cerevisia*. Tartu, 2006. 113 p.
124. **Kersti Lilleväli.** Gata3 and Gata2 in inner ear development. Tartu, 2007. 123 p.
125. **Kai Rünk.** Comparative ecology of three fern species: *Dryopteris carthusiana* (Vill.) H.P. Fuchs, *D. expansa* (C. Presl) Fraser-Jenkins & Jermy and *D. dilatata* (Hoffm.) A. Gray (Dryopteridaceae). Tartu, 2007. 143 p.
126. **Aveliina Helm.** Formation and persistence of dry grassland diversity: role of human history and landscape structure. Tartu, 2007. 89 p.
127. **Leho Tedersoo.** Ectomycorrhizal fungi: diversity and community structure in Estonia, Seychelles and Australia. Tartu, 2007. 233 p.
128. **Marko Mägi.** The habitat-related variation of reproductive performance of great tits in a deciduous-coniferous forest mosaic: looking for causes and consequences. Tartu, 2007. 135 p.
129. **Valeria Lulla.** Replication strategies and applications of Semliki Forest virus. Tartu, 2007. 109 p.
130. **Ülle Reier.** Estonian threatened vascular plant species: causes of rarity and conservation. Tartu, 2007. 79 p.
131. **Inga Jüriado.** Diversity of lichen species in Estonia: influence of regional and local factors. Tartu, 2007. 171 p.
132. **Tatjana Krama.** Mobbing behaviour in birds: costs and reciprocity based cooperation. Tartu, 2007. 112 p.
133. **Signe Saumaa.** The role of DNA mismatch repair and oxidative DNA damage defense systems in avoidance of stationary phase mutations in *Pseudomonas putida*. Tartu, 2007. 172 p.
134. **Reedik Mägi.** The linkage disequilibrium and the selection of genetic markers for association studies in european populations. Tartu, 2007. 96 p.
135. **Priit Kilgas.** Blood parameters as indicators of physiological condition and skeletal development in great tits (*Parus major*): natural variation and application in the reproductive ecology of birds. Tartu, 2007. 129 p.
136. **Anu Albert.** The role of water salinity in structuring eastern Baltic coastal fish communities. Tartu, 2007. 95 p.
137. **Kärt Padari.** Protein transduction mechanisms of transportans. Tartu, 2008. 128 p.
138. **Siiri-Lii Sandre.** Selective forces on larval colouration in a moth. Tartu, 2008. 125 p.
139. **Ülle Jõgar.** Conservation and restoration of semi-natural floodplain meadows and their rare plant species. Tartu, 2008. 99 p.
140. **Lauri Laanisto.** Macroecological approach in vegetation science: generality of ecological relationships at the global scale. Tartu, 2008. 133 p.
141. **Reidar Andreson.** Methods and software for predicting PCR failure rate in large genomes. Tartu, 2008. 105 p.
142. **Birgot Paavel.** Bio-optical properties of turbid lakes. Tartu, 2008. 175 p.

143. **Kaire Torn.** Distribution and ecology of charophytes in the Baltic Sea. Tartu, 2008, 98 p.
144. **Vladimir Vimberg.** Peptide mediated macrolide resistance. Tartu, 2008, 190 p.
145. **Daima Örd.** Studies on the stress-inducible pseudokinase TRB3, a novel inhibitor of transcription factor ATF4. Tartu, 2008, 108 p.
146. **Lauri Saag.** Taxonomic and ecologic problems in the genus *Lepraria* (*Stereocaulaceae*, lichenised *Ascomycota*). Tartu, 2008, 175 p.
147. **Ulvi Karu.** Antioxidant protection, carotenoids and coccidians in greenfinches – assessment of the costs of immune activation and mechanisms of parasite resistance in a passerine with carotenoid-based ornaments. Tartu, 2008, 124 p.
148. **Jaanus Remm.** Tree-cavities in forests: density, characteristics and occupancy by animals. Tartu, 2008, 128 p.
149. **Epp Moks.** Tapeworm parasites *Echinococcus multilocularis* and *E. granulosus* in Estonia: phylogenetic relationships and occurrence in wild carnivores and ungulates. Tartu, 2008, 82 p.
150. **Eve Eensalu.** Acclimation of stomatal structure and function in tree canopy: effect of light and CO₂ concentration. Tartu, 2008, 108 p.
151. **Janne Pullat.** Design, functionlization and application of an *in situ* synthesized oligonucleotide microarray. Tartu, 2008, 108 p.
152. **Marta Putrinš.** Responses of *Pseudomonas putida* to phenol-induced metabolic and stress signals. Tartu, 2008, 142 p.
153. **Marina Semtšenko.** Plant root behaviour: responses to neighbours and physical obstructions. Tartu, 2008, 106 p.
154. **Marge Starast.** Influence of cultivation techniques on productivity and fruit quality of some *Vaccinium* and *Rubus* taxa. Tartu, 2008, 154 p.
155. **Age Tats.** Sequence motifs influencing the efficiency of translation. Tartu, 2009, 104 p.
156. **Radi Tegova.** The role of specialized DNA polymerases in mutagenesis in *Pseudomonas putida*. Tartu, 2009, 124 p.
157. **Tsipe Aavik.** Plant species richness, composition and functional trait pattern in agricultural landscapes – the role of land use intensity and landscape structure. Tartu, 2008, 112 p.
158. **Kaja Kiiver.** Semliki forest virus based vectors and cell lines for studying the replication and interactions of alphaviruses and hepaciviruses. Tartu, 2009, 104 p.
159. **Meelis Kadaja.** Papillomavirus Replication Machinery Induces Genomic Instability in its Host Cell. Tartu, 2009, 126 p.
160. **Pille Hallast.** Human and chimpanzee Luteinizing hormone/Chorionic Gonadotropin beta (*LHB/CGB*) gene clusters: diversity and divergence of young duplicated genes. Tartu, 2009, 168 p.
161. **Ain Vellak.** Spatial and temporal aspects of plant species conservation. Tartu, 2009, 86 p.

162. **Triinu Remmel.** Body size evolution in insects with different colouration strategies: the role of predation risk. Tartu, 2009, 168 p.
163. **Jaana Salujõe.** Zooplankton as the indicator of ecological quality and fish predation in lake ecosystems. Tartu, 2009, 129 p.
164. **Ele Vahtmäe.** Mapping benthic habitat with remote sensing in optically complex coastal environments. Tartu, 2009, 109 p.
165. **Liisa Metsamaa.** Model-based assessment to improve the use of remote sensing in recognition and quantitative mapping of cyanobacteria. Tartu, 2009, 114 p.
166. **Pille Säälük.** The role of endocytosis in the protein transduction by cell-penetrating peptides. Tartu, 2009, 155 p.
167. **Lauri Peil.** Ribosome assembly factors in *Escherichia coli*. Tartu, 2009, 147 p.
168. **Lea Hallik.** Generality and specificity in light harvesting, carbon gain capacity and shade tolerance among plant functional groups. Tartu, 2009, 99 p.
169. **Mariliis Tark.** Mutagenic potential of DNA damage repair and tolerance mechanisms under starvation stress. Tartu, 2009, 191 p.
170. **Riinu Rannap.** Impacts of habitat loss and restoration on amphibian populations. Tartu, 2009, 117 p.
171. **Maarja Adojaan.** Molecular variation of HIV-1 and the use of this knowledge in vaccine development. Tartu, 2009, 95 p.
172. **Signe Altmäe.** Genomics and transcriptomics of human induced ovarian folliculogenesis. Tartu, 2010, 179 p.
173. **Triin Suvi.** Mycorrhizal fungi of native and introduced trees in the Seychelles Islands. Tartu, 2010, 107 p.
174. **Velda Lauringson.** Role of suspension feeding in a brackish-water coastal sea. Tartu, 2010, 123 p.
175. **Eero Talts.** Photosynthetic cyclic electron transport – measurement and variably proton-coupled mechanism. Tartu, 2010, 121 p.
176. **Mari Nelis.** Genetic structure of the Estonian population and genetic distance from other populations of European descent. Tartu, 2010, 97 p.
177. **Kaarel Krjutškov.** Arrayed Primer Extension-2 as a multiplex PCR-based method for nucleic acid variation analysis: method and applications. Tartu, 2010, 129 p.
178. **Egle Köster.** Morphological and genetical variation within species complexes: *Anthyllis vulneraria* s. l. and *Alchemilla vulgaris* (coll.). Tartu, 2010, 101 p.
179. **Erki Õunap.** Systematic studies on the subfamily Sterrhinae (Lepidoptera: Geometridae). Tartu, 2010, 111 p.
180. **Merike Jõesaar.** Diversity of key catabolic genes at degradation of phenol and *p*-cresol in pseudomonads. Tartu, 2010, 125 p.
181. **Kristjan Herkül.** Effects of physical disturbance and habitat-modifying species on sediment properties and benthic communities in the northern Baltic Sea. Tartu, 2010, 123 p.

182. **Arto Pulk.** Studies on bacterial ribosomes by chemical modification approaches. Tartu, 2010, 161 p.
183. **Maria Põllupüü.** Ecological relations of cladocerans in a brackish-water ecosystem. Tartu, 2010, 126 p.
184. **Toomas Silla.** Study of the segregation mechanism of the Bovine Papillomavirus Type 1. Tartu, 2010, 188 p.
185. **Gyaneshwer Chaubey.** The demographic history of India: A perspective based on genetic evidence. Tartu, 2010, 184 p.
186. **Katrin Kepp.** Genes involved in cardiovascular traits: detection of genetic variation in Estonian and Czech populations. Tartu, 2010, 164 p.
187. **Virve Sõber.** The role of biotic interactions in plant reproductive performance. Tartu, 2010, 92 p.
188. **Kersti Kangro.** The response of phytoplankton community to the changes in nutrient loading. Tartu, 2010, 144 p.
189. **Joachim M. Gerhold.** Replication and Recombination of mitochondrial DNA in Yeast. Tartu, 2010, 120 p.
190. **Helen Tammert.** Ecological role of physiological and phylogenetic diversity in aquatic bacterial communities. Tartu, 2010, 140 p.
191. **Elle Rajandu.** Factors determining plant and lichen species diversity and composition in Estonian *Calamagrostis* and *Hepatica* site type forests. Tartu, 2010, 123 p.
192. **Paula Ann Kivistik.** ColR-ColS signalling system and transposition of Tn4652 in the adaptation of *Pseudomonas putida*. Tartu, 2010, 118 p.
193. **Siim Sõber.** Blood pressure genetics: from candidate genes to genome-wide association studies. Tartu, 2011, 120 p.
194. **Kalle Kipper.** Studies on the role of helix 69 of 23S rRNA in the factor-dependent stages of translation initiation, elongation, and termination. Tartu, 2011, 178 p.
195. **Triinu Siibak.** Effect of antibiotics on ribosome assembly is indirect. Tartu, 2011, 134 p.
196. **Tambet Tõnissoo.** Identification and molecular analysis of the role of guanine nucleotide exchange factor RIC-8 in mouse development and neural function. Tartu, 2011, 110 p.
197. **Helin Räägel.** Multiple faces of cell-penetrating peptides – their intracellular trafficking, stability and endosomal escape during protein transduction. Tartu, 2011, 161 p.
198. **Andres Jaanus.** Phytoplankton in Estonian coastal waters – variability, trends and response to environmental pressures. Tartu, 2011, 157 p.
199. **Tiit Nikopensius.** Genetic predisposition to nonsyndromic orofacial clefts. Tartu, 2011, 152 p.
200. **Signe Värvi.** Studies on the mechanisms of RNA polymerase II-dependent transcription elongation. Tartu, 2011, 108 p.
201. **Kristjan Välik.** Gene expression profiling and genome-wide association studies of non-small cell lung cancer. Tartu, 2011, 98 p.

202. **Arno Põllumäe.** Spatio-temporal patterns of native and invasive zooplankton species under changing climate and eutrophication conditions. Tartu, 2011, 153 p.
203. **Egle Tammeleht.** Brown bear (*Ursus arctos*) population structure, demographic processes and variations in diet in northern Eurasia. Tartu, 2011, 143 p.
205. **Teele Jairus.** Species composition and host preference among ectomycorrhizal fungi in Australian and African ecosystems. Tartu, 2011, 106 p.
206. **Kessy Abarenkov.** PlutoF – cloud database and computing services supporting biological research. Tartu, 2011, 125 p.
207. **Marina Grigorova.** Fine-scale genetic variation of follicle-stimulating hormone beta-subunit coding gene (*FSHB*) and its association with reproductive health. Tartu, 2011, 184 p.
208. **Anu Tiitsaar.** The effects of predation risk and habitat history on butterfly communities. Tartu, 2011, 97 p.
209. **Elin Sild.** Oxidative defences in immunoeological context: validation and application of assays for nitric oxide production and oxidative burst in a wild passerine. Tartu, 2011, 105 p.
210. **Irja Saar.** The taxonomy and phylogeny of the genera *Cystoderma* and *Cystodermella* (Agaricales, Fungi). Tartu, 2011, 167 p.